

Product datasheet for MR225478L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Esrrb (NM_011934) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Esrrb (NM_011934) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Esrrb

Synonyms: Err2; Errb; Estrrb; Nr3b2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_011934 **ORF Size:** 1362 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(MR225478).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 011934.4, NP 036064.3

 RefSeq Size:
 4208 bp

 RefSeq ORF:
 1365 bp

 Locus ID:
 26380

 UniProt ID:
 Q61539

Cytogenetics: 12 40.49 cM





Gene Summary:

Transcription factor that binds a canonical ESRRB recognition (ERRE) sequence 5'TCAAGGTCA-3' localized on promoter and enhancer of targets genes regulating their expression or their transcriptional activity (PubMed:27601327, PubMed:23169531, PubMed:23508100, PubMed:26206133, PubMed:20534447, PubMed:18662995, PubMed:18957414, PubMed:27723719, PubMed:23019124). Plays a role, in a LIF-independent manner, in maintainance of self-renewal and pluripotency of embryonic and trophoblast stem cells through different signaling pathways including FGF signaling pathway and Wnt signaling pathways (PubMed:18957414, PubMed:26206133, PubMed:20534447, PubMed:23040478, PubMed:23040477, PubMed:23019124, PubMed:23169531). Upon FGF signaling pathway activation, interacts with KDM1A by directly binding to enhancer site of ELF5 and EOMES and activating their transcription leading to self-renewal of trophoblast stem cells (PubMed:26206133). Also regulates expression of multiple rod-specific genes and is required for survival of this cell type (PubMed:20534447). Plays a role as transcription factor activator of GATA6, NR0B1, POU5F1 and PERM1 (PubMed:18662995, PubMed:23508100, PubMed:18957414). Plays a role as transcription factor repressor of NFE2L2 transcriptional activity and ESR1 transcriptional activity (By similarity). During mitosis remains bound to a subset of interphase target genes, including pluripotency regulators, through the canonical ESRRB recognition (ERRE) sequence, leading to their transcriptional activation in early G1 phase (PubMed:27723719). Can coassemble on structured DNA elements with other transcription factors like SOX2, POU5F1, KDM1A and NCOA3 to trigger ESRRB-dependent gene activation (PubMed:23019124, PubMed:23169531, PubMed:18662995, PubMed:26206133). This mechanism, in the case of SOX2 corecruitment prevents the embryonic stem cells (ESCs) to epiblast stem cells (EpiSC) transition through positive regulation of NR0B1 that inhibits the EpiSC transcriptional program (PubMed:23169531). Also plays a role inner ear development by controlling expression of ion channels and transporters and in early placentation (PubMed:9285590, PubMed:17765677).[UniProtKB/Swiss-Prot Function]