

## **Product datasheet for MR225548L4V**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Odf2 (NM\_001113214) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Odf2 (NM\_001113214) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Odf2

Synonyms: AI848335; MMTEST29

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_001113214

ORF Size: 1956 bp

**ORF Nucleotide** 

•

Sequence:

The ORF insert of this clone is exactly the same as(MR225548).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 001113214.1</u>, <u>NP 001106685.1</u>

 RefSeq Size:
 2418 bp

 RefSeq ORF:
 1959 bp

 Locus ID:
 18286

 UniProt ID:
 A3KGV1

Cytogenetics: 2 20.79 cM







## **Gene Summary:**

Seems to be a major component of sperm tail outer dense fibers (ODF). ODFs are filamentous structures located on the outside of the axoneme in the midpiece and principal piece of the mammalian sperm tail and may help to maintain the passive elastic structures and elastic recoil of the sperm tail. May have a modulating influence on sperm motility. Functions as a general scaffold protein that is specifically localized at the distal/subdistal appendages of mother centrioles. Component of the centrosome matrix required for the localization of PLK1 and NIN to the centrosomes. Required for the formation and/or maintenance of normal CETN1 assembly (By similarity).[UniProtKB/Swiss-Prot Function]