

Product datasheet for MR225565L3

Irf1 (NM_001159393) Mouse Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Irf1 (NM_001159393) Mouse Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: Irf1

Synonyms: AU020929; Irf-1

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(MR225565).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF.

ACCN: NM_001159393

ORF Size: 912 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Irf1 (NM_001159393) Mouse Tagged Lenti ORF Clone - MR225565L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001159393.1</u>, <u>NP 001152865.1</u>

RefSeq Size:2128 bpRefSeq ORF:915 bpLocus ID:16362


Cytogenetics: 11 32.0 cM



Gene Summary:

Transcriptional regulator which displays a remarkable functional diversity in the regulation of cellular responses. These include the regulation of IFN and IFN-inducible genes, host response to viral and bacterial infections, regulation of many genes expressed during hematopoiesis, inflammation, immune responses and cell proliferation and differentiation, regulation of the cell cycle and induction of growth arrest and programmed cell death following DNA damage. Stimulates both innate and acquired immune responses through the activation of specific target genes and can act as a transcriptional activator and repressor regulating target genes by binding to an interferon-stimulated response element (ISRE) in their promoters. Its target genes for transcriptional activation activity are: genes involved in anti-viral response, such as IFN-alpha/beta, DDX58/RIG-I, TNFSF10/TRAIL, OAS1/2, PIAS1/GBP, EIF2AK2/PKR and RSAD2/viperin; antibacterial response, such as NOS2/INOS; anti-proliferative response, such as p53/TP53, LOX and CDKN1A; apoptosis, such as BBC3/PUMA, CASP1, CASP7 and CASP8; immune response, such as IL7, IL12A/B and IL15, PTGS2/COX2 and CYBB; DNA damage responses and DNA repair, such as POLQ/POLH; MHC class I expression, such as TAP1, PSMB9/LMP2, PSME1/PA28A, PSME2/PA28B and B2M and MHC class II expression, such as CIITA. Represses genes involved in anti-proliferative response, such as BIRC5/survivin, CCNB1, CCNE1, CDK1, CDK2 and CDK4 and in immune response, such as FOXP3, IL4, ANXA2 and TLR4. Stimulates p53/TP53-dependent transcription through enhanced recruitment of EP300 leading to increased acetylation of p53/TP53. Plays an important role in immune response directly affecting NK maturation and activity, macrophage production of IL12, Th1 development and maturation of CD8+ T-cells. Also implicated in the differentiation and maturation of dendritic cells and in the suppression of regulatory T (Treg) cells development. Acts as a tumor suppressor and plays a role not only in antagonism of tumor cell growth but also in stimulating an immune response against tumor cells.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR225565L3