

Product datasheet for MR225663L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Opa1 (NM_133752) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Opa1 (NM_133752) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Opa1

Synonyms: 1200011N24Rik; Al225888; Al847218; lilr3; mKlAA0567

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_133752

ORF Size: 2883 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR225663).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 133752.2</u>

 RefSeq Size:
 5948 bp

 RefSeq ORF:
 2883 bp

 Locus ID:
 74143

 UniProt ID:
 P58281

Cytogenetics: 16 20.65 cM







Gene Summary:

Dynamin-related GTPase that is essential for normal mitochondrial morphology by regulating the equilibrium between mitochondrial fusion and mitochondrial fission (PubMed:11847212, PubMed:24616225, PubMed:26785494, PubMed:28746876). Coexpression of isoform 1 with shorter alternative products is required for optimal activity in promoting mitochondrial fusion (By similarity). Binds lipid membranes enriched in negatively charged phospholipids, such as cardiolipin, and promotes membrane tubulation. The intrinsic GTPase activity is low, and is strongly increased by interaction with lipid membranes (By similarity). Plays a role in remodeling cristae and the release of cytochrome c during apoptosis (PubMed:16839884, PubMed:16839885). Proteolytic processing in response to intrinsic apoptotic signals may lead to disassembly of OPA1 oligomers and release of the caspase activator cytochrome C (CYCS) into the mitochondrial intermembrane space (PubMed:16839884, PubMed:16839885). Plays a role in mitochondrial genome maintenance (By similarity).[UniProtKB/Swiss-Prot Function]