

Product datasheet for **MR225751L3V**

Spast (NM_001162870) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Spast (NM_001162870) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Spast
Synonyms:	mKIAA1083; Spg4
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001162870
ORF Size:	1842 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR225751).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_001162870.1 , NP_001156342.1
RefSeq Size:	4696 bp
RefSeq ORF:	1845 bp
Locus ID:	50850
UniProt ID:	Q9QYY8
Cytogenetics:	17 E2



[View online »](#)

Gene Summary:

ATP-dependent microtubule severing protein that specifically recognizes and cuts microtubules that are polyglutamylated (PubMed:19141076 PubMed:20530212). Preferentially recognizes and acts on microtubules decorated with short polyglutamate tails: severing activity increases as the number of glutamates per tubulin rises from one to eight, but decreases beyond this glutamylation threshold (By similarity). Severing activity is not dependent on tubulin acetylation or detyrosination (By similarity). Microtubule severing promotes reorganization of cellular microtubule arrays and the release of microtubules from the centrosome following nucleation (By similarity). It is critical for the biogenesis and maintenance of complex microtubule arrays in axons, spindles and cilia (By similarity). SPAST is involved in abscission step of cytokinesis and nuclear envelope reassembly during anaphase in cooperation with the ESCRT-III complex (By similarity). Recruited at the midbody, probably by IST1, and participates in membrane fission during abscission together with the ESCRT-III complex (By similarity). Recruited to the nuclear membrane by IST1 and mediates microtubule severing, promoting nuclear envelope sealing and mitotic spindle disassembly during late anaphase (By similarity). Required for membrane traffic from the endoplasmic reticulum (ER) to the Golgi and endosome recycling (By similarity). Recruited by IST1 to endosomes and regulates early endosomal tubulation and recycling by mediating microtubule severing (By similarity). Probably plays a role in axon growth and the formation of axonal branches (PubMed:18234839).[UniProtKB/Swiss-Prot Function]