

Product datasheet for MR225770L4

Il18r1 (NM_001161842) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Il18r1 (NM_001161842) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Il18r1
Synonyms:	Il1rrp; Il18ralpha
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR225770).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	


Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.



[View online »](#)

Plasmid Map:


ACCN: NM_001161842

ORF Size: 1116 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001161842.1](#), [NP_001155314.1](#)

RefSeq Size: 1919 bp

RefSeq ORF: 1119 bp

Locus ID: 16182

Cytogenetics: 1 B

Gene Summary: Within the IL18 receptor complex, responsible for the binding of the proinflammatory cytokine IL18, but not IL1A nor IL1B. Involved in IL18-mediated IFNG synthesis from T-helper 1 (Th1) cells (By similarity). Contributes to IL18-induced cytokine production, either independently of SLC12A3, or as a complex with SLC12A3 (PubMed:26099046).
[UniProtKB/Swiss-Prot Function]