

## Product datasheet for **MR225787L3V**

### **Auts2 (NM\_177047) Mouse Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Auts2 (NM_177047) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Auts2
Synonyms:	2700063G02Rik; A730011F23Rik; D830032G16Rik
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_177047
ORF Size:	3783 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR225787).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_177047.3</a> , <a href="#">NP_796021.2</a>
RefSeq Size:	6060 bp
RefSeq ORF:	3786 bp
Locus ID:	319974
UniProt ID:	<a href="#">A0A087WPF7</a>
Cytogenetics:	5 G2



[View online »](#)

**Gene Summary:**

Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. The PRC1-like complex that contains PCGF5, RNF2, CSNK2B, RYBP and AUTS2 has decreased histone H2A ubiquitination activity, due to the phosphorylation of RNF2 by CSNK2B. As a consequence, the complex mediates transcriptional activation (By similarity). In the cytoplasm, plays a role in axon and dendrite elongation and in neuronal migration during embryonic brain development. Promotes reorganization of the actin cytoskeleton, lamellipodia formation and neurite elongation via its interaction with RAC guanine nucleotide exchange factors, which then leads to the activation of RAC1 (PubMed:25533347).[UniProtKB/Swiss-Prot Function]