

Product datasheet for MR225908L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Kat2b (NM_001190846) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Kat2b (NM_001190846) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Kat2b

Synonyms: A930006P13Rik; AI461839; AW536563; p/CAF; Pcaf

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001190846

ORF Size: 2205 bp

ORF Nucleotide

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(MR225908).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001190846.1, NP 001177775.1

RefSeq Size: 4100 bp RefSeq ORF: 2208 bp Locus ID: 18519

Cytogenetics: 17 27.86 cM







Gene Summary:

Functions as a histone acetyltransferase (HAT) to promote transcriptional activation. Has significant histone acetyltransferase activity with core histones (H3 and H4), and also with nucleosome core particles. Also acetylates non-histone proteins, such as ACLY, PLK4 and TBX5. Inhibits cell-cycle progression and counteracts the mitogenic activity of the adenoviral oncoprotein E1A. Acts as a circadian transcriptional coactivator which enhances the activity of the circadian transcriptional activators: NPAS2-ARNTL/BMAL1 and CLOCK-ARNTL/BMAL1 heterodimers. Involved in heart and limb development by mediating acetylation of TBX5, acetylation regulating nucleocytoplasmic shuttling of TBX5. Acts as a negative regulator of centrosome amplification by mediating acetylation of PLK4. Also acetylates spermidine (By similarity).[UniProtKB/Swiss-Prot Function]