

Product datasheet for MR226050L3

Nr1i2 (NM_001098404) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Nr1i2 (NM_001098404) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Nr1i2
Synonyms:	mPXR; PXR; PXR.1; PXR.2; PXR1; SXR
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR226050).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	


Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.



[View online »](#)

Plasmid Map:


ACCN: NM_001098404

ORF Size: 1170 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001098404.1](#), [NP_001091874.1](#)

RefSeq Size: 2417 bp

RefSeq ORF: 1173 bp

Locus ID: 18171

UniProt ID: [O54915](#)

Cytogenetics: 16 B3

Gene Summary: Nuclear receptor that binds and is activated by a variety of endogenous and xenobiotic compounds. Transcription factor that activates the transcription of multiple genes involved in the metabolism and secretion of potentially harmful xenobiotics, endogenous compounds and drugs. Response to specific ligands is species-specific, due to differences in the ligand-binding domain. Binds to a response element in the promoters of the CYP3A4 and ABCB1/MDR1 genes (By similarity). Activated by naturally occurring steroids such as pregnenolone and progesterone, the cholesterol metabolite 5-beta-cholestane-3-alpha,7-alpha,12-alpha-triol, synthetic glucocorticoids and antiglucocorticoids and 16-alpha-carbonitrile (PCN).[UniProtKB/Swiss-Prot Function]