

## Product datasheet for MR226079L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Gtf2i (NM\_001080748) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Gtf2i (NM\_001080748) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Gtf2

Synonyms: 6030441I21Rik; BAP-135; Diws1t; Gtfll-I; Spin; TFll-I; WBSCR6

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001080748

ORF Size: 2874 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR226079).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001080748.1</u>

 RefSeq Size:
 4335 bp

 RefSeq ORF:
 2877 bp

 Locus ID:
 14886

 UniProt ID:
 Q9ESZ8

**Cytogenetics:** 5 74.48 cM





## **Gene Summary:**

Interacts with the basal transcription machinery by coordinating the formation of a multiprotein complex at the C-FOS promoter, and linking specific signal responsive activator complexes. Promotes the formation of stable high-order complexes of SRF and PHOX1 and interacts cooperatively with PHOX1 to promote serum-inducible transcription of a reporter gene deriven by the C-FOS serum response element (SRE). Acts as a coregulator for USF1 by binding independently two promoter elements, a pyrimidine-rich initiator (Inr) and an upstream E-box (By similarity). Required for the formation of functional ARID3A DNA-binding complexes and for activation of immunoglobulin heavy-chain transcription upon B-lymphocyte activation.[UniProtKB/Swiss-Prot Function]