

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for MR226081L4V

## Gtf2i (NM\_001080749) Mouse Tagged ORF Clone Lentiviral Particle

### **Product data:**

Lentiviral Particles
Gtf2i (NM_001080749) Mouse Tagged ORF Clone Lentiviral Particle
Gtf2i
6030441I21Rik; BAP-135; Diws1t; GtfII-I; Spin; TFII-I; WBSCR6
Puromycin
pLenti-C-mGFP-P2A-Puro (PS100093)
mGFP
NM_001080749
2886 bp
The ORF insert of this clone is exactly the same as(MR226081).
The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<u>NM 001080749.1</u>
4347 bp
2889 bp
14886
5 74.48 cM



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



#### Gtf2i (NM\_001080749) Mouse Tagged ORF Clone Lentiviral Particle – MR226081L4V

Gene Summary:Interacts with the basal transcription machinery by coordinating the formation of a<br/>multiprotein complex at the C-FOS promoter, and linking specific signal responsive activator<br/>complexes. Promotes the formation of stable high-order complexes of SRF and PHOX1 and<br/>interacts cooperatively with PHOX1 to promote serum-inducible transcription of a reporter<br/>gene deriven by the C-FOS serum response element (SRE). Acts as a coregulator for USF1 by<br/>binding independently two promoter elements, a pyrimidine-rich initiator (Inr) and an<br/>upstream E-box (By similarity). Required for the formation of functional ARID3A DNA-binding<br/>complexes and for activation of immunoglobulin heavy-chain transcription upon B-<br/>lymphocyte activation.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US