

## Product datasheet for **MR226108L4V**

### **Rbpj (NM\_009035) Mouse Tagged ORF Clone Lentiviral Particle**

#### **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Rbpj (NM_009035) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Rbpj
Synonyms:	AI843960; CBF1; Iglkjrj; Iglkrsbp; RBP-J; RBP-J kappa; RBP-Jkappa; RBPjk; Rbpsuh
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_009035
ORF Size:	1578 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR226108).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_009035.4</a>
RefSeq Size:	5541 bp
RefSeq ORF:	1581 bp
Locus ID:	19664
UniProt ID:	<a href="#">P31266</a>
Cytogenetics:	5 29.37 cM



[View online »](#)

**Gene Summary:**

Transcriptional regulator that plays a central role in Notch signaling, a signaling pathway involved in cell-cell communication that regulates a broad spectrum of cell-fate determinations (PubMed:7566092). Acts as a transcriptional repressor when it is not associated with Notch proteins. When associated with some NICD product of Notch proteins (Notch intracellular domain), it acts as a transcriptional activator that activates transcription of Notch target genes. Probably represses or activates transcription via the recruitment of chromatin remodeling complexes containing histone deacetylase or histone acetylase proteins, respectively. Specifically binds to the immunoglobulin kappa-type J segment recombination signal sequence. Binds specifically to methylated DNA. Binds to the oxygen responsive element of COX4I2 and activates its transcription under hypoxia conditions (4% oxygen) (By similarity). Negatively regulates the phagocyte oxidative burst in response to bacterial infection by repressing transcription of NADPH oxidase subunits (PubMed:26194095).[UniProtKB/Swiss-Prot Function]