

## Product datasheet for MR226130L4

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

# Ntrk2 (NM\_001025074) Mouse Tagged Lenti ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

Product Name: Ntrk2 (NM\_001025074) Mouse Tagged Lenti ORF Clone

Tag: mGFP Symbol: Ntrk2

**Synonyms:** GP145-TrkB/GP95-TrkB; Tkrb; trk-B; trkB

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

E. coli Selection: Chloramphenicol (34 ug/mL)

**ORF Nucleotide** The ORF insert of this clone is exactly the same as(MR226130).

Sequence:

Restriction Sites: Sgfl-Mlul

**Cloning Scheme:** 






<sup>\*</sup> The last codon before the Stop codon of the ORF.





#### Plasmid Map:



**ACCN:** NM\_001025074

ORF Size: 2463 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 001025074.1, NP 001020245.1

RefSeq Size: 4614 bp RefSeq ORF: 2466 bp



### Ntrk2 (NM\_001025074) Mouse Tagged Lenti ORF Clone - MR226130L4

 Locus ID:
 18212

 UniProt ID:
 P15209

 Cytogenetics:
 13 31.2 cM

Gene Summary: Receptor tyrosine kinase involved in the developme

Receptor tyrosine kinase involved in the development and the maturation of the central and the peripheral nervous systems through regulation of neuron survival, proliferation, migration, differentiation, and synapse formation and plasticity. Receptor for BDNF/brainderived neurotrophic factor and NTF4/neurotrophin-4. Alternatively can also bind NTF3/neurotrophin-3 which is less efficient in activating the receptor but regulates neuron survival through NTRK2. Upon ligand-binding, undergoes homodimerization, autophosphorylation and activation. Recruits, phosphorylates and/or activates several downstream effectors including SHC1, FRS2, SH2B1, SH2B2 and PLCG1 that regulate distinct overlapping signaling cascades. Through SHC1, FRS2, SH2B1, SH2B2 activates the GRB2-Ras-MAPK cascade that regulates for instance neuronal differentiation including neurite outgrowth. Through the same effectors controls the Ras-PI3 kinase-AKT1 signaling cascade that mainly regulates growth and survival. Through PLCG1 and the downstream protein kinase C-regulated pathways controls synaptic plasticity. Thereby, plays a role in learning and memory by regulating both short term synaptic function and long-term potentiation. PLCG1 also leads to NF-Kappa-B activation and the transcription of genes involved in cell survival. Hence, it is able to suppress anoikis, the apoptosis resulting from loss of cell-matrix interactions. Isoform GP95-TRKB may also play a role in neutrophin-dependent calcium signaling in glial cells and mediate communication between neurons and glia. [UniProtKB/Swiss-Prot Function]