

Product datasheet for **MR226161L1V**

DII1 (NM_007865) Mouse Tagged ORF Clone Lentiviral Particle

Product data:


Product Type:	Lentiviral Particles
Product Name:	DII1 (NM_007865) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	DII1
Synonyms:	Delta1
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_007865
ORF Size:	2166 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR226161).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_007865.3 , NP_031891.2
RefSeq Size:	3444 bp
RefSeq ORF:	2169 bp
Locus ID:	13388
UniProt ID:	Q61483
Cytogenetics:	17 8.95 cM
Gene Summary:	Transmembrane ligand protein of NOTCH1, NOTCH2 and NOTCH3 receptors that binds the extracellular domain (ECD) of Notch receptor in a cis and trans fashion manner (PubMed:21985982, PubMed:10958687). Following transinteraction, ligand cells produce


[View online »](#)

mechanical force that depends of a clathrin-mediated endocytosis, requiring ligand ubiquitination, EPN1 interaction, and actin polymerisation; these events promote Notch receptor extracellular domain (NECD) transendocytosis and triggers Notch signaling through induction of cleavage, hyperphosphorylation, and nuclear accumulation of the intracellular domain of Notch receptors (NICD) (PubMed:10958687, PubMed:18676613). Is required for embryonic development and maintenance of adult stem cells in many different tissues and immune system; the DLL1-induced Notch signaling is mediated through an intercellular communication that regulates cell lineage, cell specification, cell patterning and morphogenesis through effects on differentiation and proliferation (PubMed:17194759, PubMed:19562077, PubMed:18997111, PubMed:23695674, PubMed:16495313, PubMed:21238454, PubMed:22282195, PubMed:7671806, PubMed:17960184, PubMed:22529374, PubMed:19389377, PubMed:23699523, PubMed:19144989, PubMed:23688253, PubMed:23806616, PubMed:26114479, PubMed:22940113, PubMed:25220152, PubMed:20081190, PubMed:21572390, PubMed:22096075). Plays a role in brain development at different level, namely by regulating neuronal differentiation of neural precursor cells via cell-cell interaction, most likely through the lateral inhibitory system in an endogenous level dependent-manner (PubMed:7671806, PubMed:18997111). During neocortex development, Dll1-Notch signaling transmission is mediated by dynamic interactions between intermediate neurogenic progenitors and radial glia; the cell-cell interactions are mediated via dynamic and transient elongation processes, likely to reactivate/maintain Notch activity in neighboring progenitors, and coordinate progenitor cell division and differentiation across radial and zonal boundaries (PubMed:23699523). During cerebellar development, regulates Bergmann glial monolayer formation and its morphological maturation through a Notch signaling pathway (PubMed:23688253). At the retina and spinal cord level, regulates neurogenesis by preventing the premature differentiation of neural progenitors and also by maintaining progenitors in spinal cord through Notch signaling pathway (PubMed:19389377, PubMed:26114479). Also controls neurogenesis of the neural tube in a progenitor domain-specific fashion along the dorsoventral axis (PubMed:20081190). Maintains quiescence of neural stem cells and plays a role as a fate determinant that segregates asymmetrically to one daughter cell during neural stem cells mitosis, resulting in neuronal differentiation in Dll1-inheriting cell (PubMed:23695674). Plays a role in immune system development, namely the development of all T-cells and marginal zone (MZ) B cells (PubMed:15146182, PubMed:19217325). Blocks the differentiation of progenitor cells into the B-cell lineage while promoting the emergence of a population of cells with the characteristics of a T-cell/NK-cell precursor (By similarity). Upon MMP14 cleavage, negatively regulates Notch signaling in haematopoietic progenitor cells to specifically maintain normal B-cell development in bone marrow (PubMed:21572390). Also plays a role during muscle development. During early development, inhibits myoblasts differentiation from the medial dermomyotomal lip and later regulates progenitor cell differentiation (PubMed:17194759). Directly modulates cell adhesion and basal lamina formation in satellite cells through Notch signaling. Maintains myogenic progenitors pool by suppressing differentiation through down-regulation of MYOD1 and is required for satellite cell homing and PAX7 expression (PubMed:22940113). During craniofacial and trunk myogenesis suppresses differentiation of cranial mesoderm-derived and somite-derived

muscle via MYOD1 regulation but in cranial mes

Product images:



[MR226161L1] was used to prepare Lentiviral particles using [TR30037] packaging kit. HEK293T cells were transduced with MR226161L1V particle to overexpress human DII1-Myc-DDK fusion protein.