

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR226166L3V

Slc1a2 (NM_001077514) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Slc1a2 (NM_001077514) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Slc1a2
Synonyms:	1700091C19Rik; 2900019G14Rik; Al159670; Eaat2; GLT-1; GLT1; MGLT1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001077514
ORF Size:	1716 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR226166).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001077514.3</u>
RefSeq Size:	11571 bp
RefSeq ORF:	1719 bp
Locus ID:	20511
UniProt ID:	<u>P43006</u>
Cytogenetics:	2 54.13 cM



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Slc1a2 (NM_001077514) Mouse Tagged ORF Clone Lentiviral Particle – MR226166L3V
Gene Summary:	Sodium-dependent, high-affinity amino acid transporter that mediates the uptake of L- glutamate and also L-aspartate and D-aspartate (PubMed:7698742, PubMed:7557442, PubMed:9373176) Eurotions as a symporter that transports one amino acid molecule
	together with two or three Na(+) ions and one proton, in parallel with the counter-transport of one K(+) ion. Mediates Cl(-) flux that is not coupled to amino acid transport; this avoids the accumulation of negative charges due to aspartate and Na(+) symport (By similarity).
	Essential for the rapid removal of released glutamate from the synaptic cleft, and for terminating the postsynaptic action of glutamate (PubMed:9180080).[UniProtKB/Swiss-Prot

Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US