

## Product datasheet for **MR226206L2V**

### Ngfr (NM\_033217) Mouse Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Ngfr (NM_033217) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Ngfr
Synonyms:	LNGFR; p75; p75NGFR; p75NTR; Tnfrsf16
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_033217
ORF Size:	1284 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR226206).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_033217.3</a> , <a href="#">NP_150086.2</a>
RefSeq Size:	3409 bp
RefSeq ORF:	1284 bp
Locus ID:	18053
UniProt ID:	<a href="#">Q9Z0W1</a>
Cytogenetics:	11 59.01 cM



[View online »](#)

**Gene Summary:**

Low affinity neurotrophin receptor which can bind to mature NGF, BDNF, NTF3, and NTF4 (PubMed:11559852, PubMed:1317267). Forms a heterodimeric receptor with SORCS2 that binds the precursor forms of NGF (proNGF), BDNF (proBDNF) and NTF3 (proNTF3) with high affinity, and has much lower affinity for mature NGF and BDNF (PubMed:22155786, PubMed:24908487, PubMed:27457814). Plays an important role in differentiation and survival of specific neuronal populations during development (PubMed:1317267, PubMed:11559852). Can mediate cell survival as well as cell death of neural cells (PubMed:1317267, PubMed:11559852, PubMed:24908487). The heterodimeric receptor formed with SORCS2 plays a role in proBDNF-dependent synaptic plasticity, in hippocampal long term depression (LTD) and long term potentiation (LTP) (PubMed:27457814). Plays a role in the inactivation of RHOA (By similarity). Plays a role in the regulation of the translocation of GLUT4 to the cell surface in adipocytes and skeletal muscle cells in response to insulin, probably by regulating RAB31 activity, and thereby contributes to the regulation of insulin-dependent glucose uptake (PubMed:22460790). Necessary for the circadian oscillation of the clock genes ARNTL/BMAL1, PER1, PER2 and NR1D1 in the suprachiasmatic nucleus (SCN) of the brain and in liver and of the genes involved in glucose and lipid metabolism in the liver (PubMed:23785138).[UniProtKB/Swiss-Prot Function]