

Product datasheet for MR226304L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Kcnk18 (NM_207261) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Kcnk18 (NM 207261) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Kcnk18

Synonyms: Gm781; Tresk; Tresk-2; Trik

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_207261

 ORF Size:
 1182 bp

ORF Nucleotide

OTI Disclaimer:

The

Sequence:

The ORF insert of this clone is exactly the same as(MR226304).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 207261.3, NP 997144.1

 RefSeq Size:
 3032 bp

 RefSeq ORF:
 1185 bp

 Locus ID:
 332396

 UniProt ID:
 Q6VV64

 Cytogenetics:
 19 D3







Gene Summary:

Outward rectifying potassium channel. Produces rapidly activating outward rectifier K(+) currents. May function as background potassium channel that sets the resting membrane potential. Channel activity is directly activated by calcium signal. Activated by the G(q)-protein coupled receptor pathway. The calcium signal robustly activates the channel via calcineurin, whereas the anchoring of 14-3-3/YWHAH interferes with the return of the current to the resting state after activation. Inhibited also by arachidonic acid and other naturally occurring unsaturated free fatty acids. Channel activity is also enhanced by volatile anesthetics, such as isoflurane. Appears to be the primary target of hydroxy-alpha-sanshool, an ingredient of Schezuan pepper. May be involved in the somatosensory function with special respect to pain sensation.[UniProtKB/Swiss-Prot Function]