

Product datasheet for **MR226311**

Prkg1 (NM_011160) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Prkg1 (NM_011160) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Prkg1
Synonyms:	AW125416; CGKI; Gm19690; Prkg1b; Prkgr1b
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>MR226311 representing NM_011160
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGGCACCTGCGGGATTTACAGTATGCGCTCCAGGAGAAGATCGAGGAGCTGAGGCAGCGGGATGCTC
 TCATCGATGAGCTGGAGCTGGAGTTGGATCAGAAGGACGAACTGATCCAGAAGCTGCAGAACGAACTGGA
 CAAGTATCGTCCGGTATCCGGCCGGCCACCCAGCAGGCCGAGAAGCAGAGCGCCAGCACCTTGCAGGGT
 GAACCGCGCACCAAGCGCCAGGCGATCTCCGCGGAGCCACCGCCTTCGACATCCAGGATCTCAGCCACG
 TGACCCTGCCCTTACCCCAAGAGCCACAGTCAAGGATCTCATAAAGGAGGCTATCCTTGACAAATGA
 CTTTATGAAGAACTTGGAGCTGTACAGATCCAGGAGATTGTGGACTGTATGTACCCCGTGAATACGGC
 AAGGACAGCTGCATCATCAAGGAAGGCGATGTGGGGTCACTGGTGTACGTATGGAAGATGGGAAGTTG
 AAGTACAAAAAGGCGTGAAGCTCTGCACCATGGGTCTGGAAAAGTGTTCGGGGAGCTGGCTATACT
 TTACAACGTACCCGGACAGCGACCGTCAAGACTCTTGTAAATGTGAAACTCTGGCCATCGATCGACAA
 TGTTTTCAAACAATAATGATGAGGACAGGACTCATCAAGCATACCGAGTACATGGAATTTTTAAAAAGTG
 TTCCAACATTCAGAGCCTTCTGATGAAATCCTCAGCAAGCTGGCTGATGTCCTCGAAGAGACCCACTA
 TGAAAAATGGAGAAATATATCATCAGGCAGGGTGCAGAGGAGACACCTTCTTATCATCAGTAAAGGGCAG
 GTGAATGTTACTCGAGAAGACTACCAAGTGAAGACCCAGTCTTCTTAGAACTTTAGGGAAGGGAGATT
 GGTTTGGAGAGAAAGCGTTGCAGGGGGAGGATGTGAGAACAGCAAATGTTATTGCCCGAGAAGCTGTCAC
 CTGCCCTTGATTGACAGAGACTCTTCAAGCATTTGATTGGAGGACTGGACGATGTTTTCAACAAAGCA
 TATGAGGACGCAAGCAAAAGCAAAATATGAAGCCGAAGCTGCCTTCTCGCCAACTGAAGCTGTCTG
 ATTTCAACATCATTGACACCCCTTGAGTTGGAGTTTCGACGAGTAGAGCTGGTCCAGTTGAAAAGTGA
 AGAATCCAAAACATTTGCAATGAAGATCCTCAAGAAACGCCACATTGTGGACACCAGACAGCAGGAGCAC
 ATCCGCTCAGAGAAGCAGATCATGCAGGGGCTCATTCTGACTTCATTGTGAGGCTGTACAGGACATTTA
 AAGACAGCAAATACTTGTATATGTTGATGGAAGCGTGCCTGGGTGGAGAGCTCTGGACTATTCTCAGGGA
 TAGGGGTCGTTTGAAGACTCAACAACCAGTTTTACTGTCATGTGTGGTAGAAGCATTTCGCTATCTG
 CATTCAAAGGAATCATTACAGGGACCTCAAGCCGAGAATCTCATCCTAGATCATCGAGGCTATGCCA
 AACTGGTTGACTTTGGCTTTGCAAAGAAAATAGGATTTGGAAAGAAAACATGGACTTTTTGTGGGACTCC
 AGAATATGTAGCCCCAGAGATCATCCTGAACAAAGGCCATGACATTTAGCTGACTACTGGTCACTAGGA
 ATTCTGATGTATGAGCTTCTGACTGGCAGCCACCTTTCTCAGGCCAGATCCAATGAAAACCTACAATA
 TCATACTGCGGGGATTGACATGATAGAATTTCAAAGAAGATTGCAAAAATGCTGCTAATTTAATTA
 AAAACTATGCAGGGACAACCCATCAGAGAGTTAGGAAATTTGAAAACGGAGTCAAAGACATTCAGAAA
 CACAAGTGGTTTGGGGCTTTAATTGGGAAGGCTTAAGAAAAGGCACCTTGACACCTCCCATAATTTCAA
 GTGTTGCGTCACCCACAGACACAAGCAATTTTGACAGTTCCCTGAGGACAGCGATGAGCCACCACCTGA
 TGACAACCTCAGGCTGGGACATAGACTTC

ACGCGTACGCGGGCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR226311 representing NM_011160
Red=Cloning site Green=Tags(s)

```

MGTLRDLQYALQEIEELRQRDALIDELELELDQKDELIQKLQNELDKYRSVIRPATQQAQKQSASTLQG
EPRTKRQAISAEPTAFDIQDLSHVTLPFYPKSPQSKDLIKEAILDNDFMKNLELSQIQEIVDCMYPVEYG
KDSCIIKEGDVGLVYVMEDGKVEVTKEGVKLC TMGPGKVF GELAILYNCTRTATVKTLVNVKLWAIDRQ
CFQTIMMRTGLIKHTEYMEFLKSVPTFQSLPDEILSKLADVLEETHYENGEYIIRQGARGDTFFIISKGQ
VNVTRDSPSEDVFLRTLGGKDWFGKALQGEDVRTANVIAAEAVTCLVIDRDSFKHLIGGLDDVSNKA
YEDAEAKAKYEAEEAFFANLKLSDFNIDTLGVGGFGRVELVQLKSEESKTFAMKILKKRHIVDTRQOEH
IRSEKQIMQGAHSDFIVRLYRTFKDSKYLMLMEACLGGELWTILRDRGSFEDSTTRFYTACVVEAFAYL
HSKGIIYRDLKPENLILDHRGYAKLVDFGFAKKIGFGKKTWTFCGTPEYVAPEIILNKGHDSADYWSLG
ILMYELLTGSPFSGPDPMKTYNIIILRGIDMIEFPKKIAKNAANLIKCLRDNPSERLGNLKNGVKDIQK
HKWFEGFNWEGLRKGTLPPIIPSVASPTDTSNFDSPEDSDEPPPPDDNSGWDIDF
    
```

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mm9009_g11.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_011160

ORF Size: 2058 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_011160.3</u> , <u>NP_035290.1</u>
RefSeq Size:	6915 bp
RefSeq ORF:	2061 bp
Locus ID:	19091
UniProt ID:	<u>P0C605</u>
Cytogenetics:	19 C1
MW:	78.2 kDa

Gene Summary:

Serine/threonine protein kinase that acts as key mediator of the nitric oxide (NO)/cGMP signaling pathway. GMP binding activates PRKG1, which phosphorylates serines and threonines on many cellular proteins. Numerous protein targets for PRKG1 phosphorylation are implicated in modulating cellular calcium, but the contribution of each of these targets may vary substantially among cell types. Proteins that are phosphorylated by PRKG1 regulate platelet activation and adhesion, smooth muscle contraction, cardiac function, gene expression, feedback of the NO-signaling pathway, and other processes involved in several aspects of the CNS like axon guidance, hippocampal and cerebellar learning, circadian rhythm and nociception. Smooth muscle relaxation is mediated through lowering of intracellular free calcium, by desensitization of contractile proteins to calcium, and by decrease in the contractile state of smooth muscle or in platelet activation. Regulates intracellular calcium levels via several pathways: phosphorylates MRV1/IRAG and inhibits IP3-induced Ca(2+) release from intracellular stores, phosphorylation of KCNMA1 (BKCa) channels decreases intracellular Ca(2+) levels, which leads to increased opening of this channel. PRKG1 phosphorylates the canonical transient receptor potential channel (TRPC) family which inactivates the associated inward calcium current. Another mode of action of NO/cGMP/PKG1 signaling involves PKGI-mediated inactivation of the Ras homolog gene family member A (RhoA). Phosphorylation of RHOA by PRKG1 blocks the action of this protein in myriad processes: regulation of RHOA translocation; decreasing contraction; controlling vesicle trafficking, reduction of myosin light chain phosphorylation resulting in vasorelaxation. Activation of PRKG1 by NO signaling alters also gene expression in a number of tissues. In smooth muscle cells, increased cGMP and PRKG1 activity influence expression of smooth muscle-specific contractile proteins, levels of proteins in the NO/cGMP signaling pathway, down-regulation of the matrix proteins osteopontin and thrombospondin-1 to limit smooth muscle cell migration and phenotype. Regulates vasodilator-stimulated phosphoprotein (VASP) functions in platelets and smooth muscle.[UniProtKB/Swiss-Prot Function]

Product images:


Circular map for MR226311