

Product datasheet for MR226383L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Dclk2 (NM_001195498) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Dclk2 (NM_001195498) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Dclk2

Synonyms: 6330415M09Rik; AU044875; CL2; Clic; Click-II; CLICK2; Dcamk; Dcamkl2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001195498

ORF Size: 2145 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR226383).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001195498.1, NP 001182427.1

RefSeq Size: 4108 bp
RefSeq ORF: 2148 bp
Locus ID: 70762
UniProt ID: Q6PGN3

Cytogenetics: 3 F1







Gene Summary:

This gene encodes a member of the protein kinase superfamily and the doublecortin family. The protein encoded by this gene contains two N-terminal doublecortin domains, which bind microtubules and regulate microtubule polymerization, a C-terminal serine/threonine protein kinase domain, which shows substantial homology to Ca2+/calmoduline-dependent protein kinase, and a serine/proline-rich domain in between the doublecortin and the protein kinase domains, which mediates multiple protein-protein interactions. The microtubule-polymerizing activity of the encoded protein is independent of its protein kinase activity. This gene and the DCX gene, another family member, share function in the establishment of hippocampal organization and their absence results in a severe epileptic phenotype and lethality, as described in human patients with lissencephaly. Multiple alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Sep 2010]