

## Product datasheet for MR226393L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Srsf7 (NM\_001195486) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Srsf7 (NM\_001195486) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Srsf7

**Synonyms:** 9G8; 35kDa; 9430065L19Rik; NX-9; NX-96; Sf; Sfrs7

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_001195486

ORF Size: 681 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR226393).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 001195486.1</u>, <u>NP 001182415.1</u>

 RefSeq Size:
 2268 bp

 RefSeq ORF:
 684 bp

 Locus ID:
 225027

 UniProt ID:
 Q8BL97

 Cytogenetics:
 17 E3







## **Gene Summary:**

The protein encoded by this gene is a member of the serine/arginine (SR)-rich family of pre-mRNA splicing factors, which constitute part of the spliceosome. Each of these factors contains an RNA recognition motif (RRM) for binding RNA and an RS domain for binding other proteins. The RS domain is rich in serine and arginine residues and facilitates interaction between different SR splicing factors. In addition to being critical for mRNA splicing, the SR proteins have also been shown to be involved in mRNA export from the nucleus and in translation. Five transcript variants, four of them protein-coding and the other not protein-coding, have been found for this gene. [provided by RefSeq, Sep 2010]