

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR226419L4V

Dhx36 (NM_028136) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Dhx36 (NM_028136) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Dhx36
Synonyms:	2810407E23Rik; Al452301; AU022184; Ddx36; mKlAA1488
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_028136
ORF Size:	3003 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR226419).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 028136.2, NP 082412.2</u>
RefSeq Size:	4975 bp
RefSeq ORF:	3006 bp
Locus ID:	72162
UniProt ID:	<u>Q8VHK9</u>
Cytogenetics:	3 E1



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Gene Summary:

Multifunctional ATP-dependent helicase that unwinds G-guadruplex (G4) structures (PubMed:25611385). Plays a role in many biological processes such as genomic integrity, gene expression regulations and as a sensor to initiate antiviral responses (PubMed:21703541, PubMed:21590736). G4 structures correspond to helical structures containing guanine tetrads (By similarity). Binds with high affinity to and unwinds G4 structures that are formed in nucleic acids (G4-ADN and G4-RNA) (By similarity). Plays a role in genomic integrity (By similarity). Converts the G4-RNA structure present in telomerase RNA template component (TREC) into a double-stranded RNA to promote P1 helix formation that acts as a template boundary ensuring accurate reverse transcription (By similarity). Plays a role in transcriptional regulation. Resolves G4-DNA structures in promoters of genes, such as YY1, KIT/c-kit and ALPL and positively regulates their expression (PubMed:25611385) (By similarity). Plays a role in post-transcriptional regulation (By similarity). Unwinds a G4-RNA structure located in the 3' UTR polyadenylation site of the pre-mRNA TP53 and stimulates TP53 pre-mRNA 3'-end processing in response to ultraviolet (UV)-induced DNA damage (By similarity). Binds to the precursor-microRNA-134 (pre-miR-134) terminal loop and regulates its transport into the synapto-dendritic compartment (By similarity). Involved in the pre-miR-134dependent inhibition of target gene expression and the control of dendritic spine size (By similarity). Plays a role in the regulation of cytoplasmic mRNA translation and mRNA stability (By similarity). Binds to both G4-RNA structures and alternative non-quadruplex-forming sequence within the 3' UTR of the PITX1 mRNA regulating negatively PITX1 protein expression (By similarity). Binds to both G4-RNA structure in the 5'-UTR and AU-rich elements (AREs) localized in the 3' UTR of NKX2-5 mRNA to either stimulate protein translation or induce mRNA decay in an ELAVL1-dependent manner, respectively (By similarity). Binds also to ARE sequences present in several mRNAs mediating exosome-mediated 3'-5' mRNA degradation (By similarity). Involved in cytoplasmic urokinase-type plasminogen activator (uPA) mRNA decay (By similarity). Component of a multi-helicase-TICAM1 complex that acts as a cytoplasmic sensor of viral double-stranded RNA (dsRNA) and plays a role in the activation of a cascade of antiviral responses including the induction of proinflammatory cytokines via the adapter molecule TICAM1 (PubMed:21703541). Required for the early embryonic development and hematopoiesis (PubMed:22422825). Involved in the regulation of cardioblast differentiation and proliferation during heart development (PubMed:26489465). Involved in spermatogonia differentiation (PubMed:25611385). May play a role in ossification (PubMed:21590736).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US