

## Product datasheet for MR226455L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Ddx4 (NM\_001145885) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Ddx4 (NM\_001145885) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Ddx4

Synonyms: AV206478; Mvh; VASA

Mammalian Cell

Selection:

Puromycin

Vector:

pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_001145885

ORF Size: 2187 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(MR226455).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** NM 001145885.1, NP 001139357.1

RefSeq Size: 2850 bp RefSeq ORF: 2187 bp Locus ID: 13206

Cytogenetics: 13 63.87 cM





## **Gene Summary:**

ATP-dependent RNA helicase required during spermatogenesis to repress transposable elements and preventing their mobilization, which is essential for the germline integrity (PubMed:20439430, PubMed:28633017). Acts via the piRNA metabolic process, which mediates the repression of transposable elements during meiosis by forming complexes composed of piRNAs and Piwi proteins and governs the methylation and subsequent repression of transposons (PubMed:20439430, PubMed:28633017). Involved in the secondary piRNAs metabolic process, the production of piRNAs in fetal male germ cells through a pingpong amplification cycle (PubMed:20439430, PubMed:28633017). Required for PIWIL2 slicing-triggered piRNA biogenesis: helicase activity enables utilization of one of the slice cleavage fragments generated by PIWIL2 and processing these pre-piRNAs into piRNAs (PubMed:28633017). [UniProtKB/Swiss-Prot Function]