

Product datasheet for **MR226534L3V**

Btrc (NM_001037758) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Btrc (NM_001037758) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Btrc
Synonyms:	b-TrCP; beta-TrCP; Beta-Trcp1; E3RS-IkappaB; E3RSIkappaB; Fbw1a; FWD1; HOS; Slimb
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001037758
ORF Size:	1815 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR226534).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_001037758.2 , NP_001032847.2
RefSeq Size:	6358 bp
RefSeq ORF:	1818 bp
Locus ID:	12234
UniProt ID:	Q3ULA2
Cytogenetics:	19 C3


[View online »](#)

Gene Summary:

Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins. Recognizes and binds to phosphorylated target proteins (PubMed:10097128, PubMed:16371461, PubMed:18782782, PubMed:9859996, PubMed:9990853, PubMed:21911472). SCF(BTRC) mediates the ubiquitination of phosphorylated NFKB, ATF4, CDC25A, DLG1, FBXO5, PER1, SMAD3, SMAD4, SNAI1 and probably NFKB2. SCF(BTRC) mediates the ubiquitination of CTNNB1 and participates in Wnt signaling (By similarity). SCF(BTRC) mediates the ubiquitination of NFKBIA, NFKBIB and NFKBIE; the degradation frees the associated NFKB1 to translocate into the nucleus and to activate transcription (PubMed:9859996, PubMed:10097128). Ubiquitination of NFKBIA occurs at 'Lys-21' and 'Lys-22' (PubMed:9859996, PubMed:10097128). SCF(BTRC) mediates the ubiquitination of CEP68; this is required for centriole separation during mitosis (By similarity). SCF(BTRC) mediates the ubiquitination and subsequent degradation of nuclear NFE2L1 (PubMed:21911472). Has an essential role in the control of the clock-dependent transcription via degradation of phosphorylated PER1 and PER2 (PubMed:18782782). May be involved in ubiquitination and subsequent proteasomal degradation through a DBB1-CUL4 E3 ubiquitin-protein ligase (By similarity). Required for activation of NFKB-mediated transcription by IL1B, MAP3K14, MAP3K1, IKBKB and TNF (By similarity). Required for proteolytic processing of GLI3 (PubMed:16371461). Mediates ubiquitination of REST, thereby leading to its proteasomal degradation (By similarity).[UniProtKB/Swiss-Prot Function]