

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR226671L4V

Htr2c (NM_008312) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Lentiviral Particles
Htr2c (NM_008312) Mouse Tagged ORF Clone Lentiviral Particle
Htr2c
5-HT2C; 5-HT2cR; 5-HTR2C; 5HT1c; Htr1; Htr1c; S; SR1
Puromycin
pLenti-C-mGFP-P2A-Puro (PS100093)
mGFP
NM_008312
1377 bp
The ORF insert of this clone is exactly the same as(MR226671).
The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<u>NM 008312.4, NP 032338.3</u>
4765 bp
1380 bp
15560
<u>P34968</u>
X 68.46 cM



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Htr2c (NM_008312) Mouse Tagged ORF Clone Lentiviral Particle – MR226671L4V

Gene Summary:Serotonin (5-hydroxytryptamine, 5-HT), a neurotransmitter, elicits a wide array of
physiological effects by binding to several receptor subtypes, including the 5-HT2 family of
seven-transmembrane-spanning, G-protein-coupled receptors, which activate phospholipase
C and D signaling pathways. This gene encodes the 2C subtype of serotonin receptor and its
mRNA is subject to multiple RNA editing events, where genomically encoded adenosine
residues are converted to inosines. RNA editing is predicted to alter amino acids within the
second intracellular loop of the 5-HT2C receptor and generate receptor isoforms that differ in
their ability to interact with G proteins and the activation of phospholipase C and D signaling
cascades, thus modulating serotonergic neurotransmission in the central nervous system.
Studies in rodents show altered patterns of RNA editing in response to drug treatments and
stressful situations. [provided by RefSeq, Jul 2008]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US