

## Product datasheet for **MR226734**

### Vav3 (NM\_146139) Mouse Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Vav3 (NM_146139) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Vav3
Synonyms:	A530094I06Rik; AA986410; Idd18.1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR226734 representing NM_146139 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGCCAATTTTTACCTTTGTTTCAGAACAAGGGCCATTCAAACCACCAGAGAAACGGACCAATGGACTTC  
GGAGAGCTTCCAGACAGGTGGACCCAGGTTTGCCGAAGATGCAGGTCATTAGGAACACTGGCACCCC  
GGCCCCAGGGCTTCATGAAGGACCACCATTGCATATCCAGGCAGGCACACTGTTGAACTTCTGAGAGGA  
GATGCACACAGTGTGTTCTGGCAGGGACGAACTTAGCTTCAGGAGAGGTTGGATTTTTCCCAAGTGATG  
CTGTCAAACCCTCTCCATGTGTACCTAAACCAGTAGATTATTCTTGCCAACCCTGGTATGCTGGCCCAAT  
GGAAAGATTGCAGGCAGAACTGAACTTATTAACAGGGTGAATAGTACTTACCTTGAAGGCACAGGACC  
AAAGAATCTGGAGAATATGCAATTAGCATTAAAGTACAATAATGAAGCAAAACACATCAAGATTTTAAACA  
GAGATGGCTTTTTTACATTGCAGAAAAAGAAAATTTAAAAGCTTAATGGAACCTGTGGAGTACTACAA  
GCACCATCTCTCAAAGAAGGGTTCAGGACCTTAGATACAACGCTACAGTTTCCATACAAGGAGCCAGAA  
CAGCCAGCTGGACAGAGGGTAATAGAACAGGCAACAGCTTGTGAGTCCCAAAGTACTGGGTATTGCCA  
TCGCTCGGTACGACTTCTGTGCAAGAGATATGCGGGAGCTCTCCCTATTGAAAGGAGACATGGTGAAGAT  
TTACACGAAGATGAGTGCAAATGGCTGGTGGAGAGGTGAAGTAAATGGCAGGGTGGGCTGGTTTCCATCC  
ACATATGTGGAAGAAGATGAA

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

**Protein Sequence:** >MR226734 representing NM\_146139  
Red=Cloning site Green=Tags(s)


MPIFTFVSEQGPFPKPEKRTNGLRRASRQVDPGLPKMQVIRNYTGTAPAGLHEGPPLHIQAGDTEVLLRG  
 DAHSVFWQGRNLASGEVGFPPSDAVKPSVCPKVPDYSCQPWYAGPMERLQAETELINRVNSTYLVRHRT  
 KESGEYATSIKYNNEAKHIKILTRDGFHIAENRKFKSLMELVEYKHHSLKEGFRTLDTLQFPYKEPE  
 QPAGQRGNRTGNSLLSPKVLGIAIARYDFCARDMRELSLLKGDVMKIYTKMSANGWWRGEVNGRNVGFPS  
 TYVEEDE

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Chromatograms:** [https://cdn.origene.com/chromatograms/mm9038\\_f11.zip](https://cdn.origene.com/chromatograms/mm9038_f11.zip)

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**ACCN:** NM\_146139

**ORF Size:** 861 bp


**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_146139.2</a> , <a href="#">NP_666251.1</a>
<b>RefSeq Size:</b>	3061 bp
<b>RefSeq ORF:</b>	864 bp
<b>Locus ID:</b>	57257
<b>UniProt ID:</b>	<a href="#">Q9R0C8</a>
<b>Cytogenetics:</b>	3 F3
<b>MW:</b>	33.1 kDa
<b>Gene Summary:</b>	<p>Exchange factor for GTP-binding proteins RhoA, RhoG and, to a lesser extent, Rac1. Binds physically to the nucleotide-free states of those GTPases (By similarity). Plays an important role in angiogenesis. Its recruitment by phosphorylated EPHA2 is critical for EFNA1-induced RAC1 GTPase activation and vascular endothelial cell migration and assembly. May be important for integrin-mediated signaling, at least in some cell types. In osteoclasts, along with SYK tyrosine kinase, required for signaling through integrin alpha-v/beta-1 (ITAGV-ITGB1), a crucial event for osteoclast proper cytoskeleton organization and function. This signaling pathway involves RAC1, but not RHO, activation. Necessary for proper wound healing. In the course of wound healing, required for the phagocytotic cup formation preceding macrophage phagocytosis of apoptotic neutrophils. Responsible for integrin beta-2-mediated macrophage adhesion and, to a lesser extent, contributes to beta-3-mediated adhesion. Does not affect integrin beta-1-mediated adhesion.[UniProtKB/Swiss-Prot Function]</p>

Product images:



Circular map for MR226734