

Product datasheet for **MR226792**

Nfkb1 (NM_008689) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Nfkb1 (NM_008689) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Nfkb1
Synonyms:	NF-kappaB; NF-kappaB1; NF-KB1; p50; p50/p105; p105
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>MR226792 ORF sequence
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGCAGACGATGATCCCTACGGAACGGCAAAATGTTTCATTTGAACACTGCTTTGACTCACTCAATAT
 TTAATGCAGAATTATATTCACCAGAAAATACCACGTCAACAGATGGCCCATACCTTCAAATATTAGAGCA
 ACCAAAACAGAGGGGATTTTCGATTCCGCTATGTGTGTGAAGGCCCATCACACGGAGGGCTTCCGGGAGCC
 TCTAGTGAGAAGAACAAGAAATCCTACCACAGGTCAAAATTTGCAACTATGTGGGGCTGCAAAGGTTA
 TCGTTTCAGTTGGTCACAAATGGAAAAACATCCACCTGCACGCCACAGCCTGGTGGGCAAGCACTGTGA
 GGACGGGGTATGCACCGTAACAGCAGGACCCAAGGACATGGTGGTTGGCTTTGCAAACCTGGGAATACTT
 CATGTGACTAAGAAAAAGGTATTTGAAACTGGAAGCACGGATGACAGAGCGTGTATTAGGGGTATA
 ATCCTGGACTTCTGGTGCATTCTGACCTTGCCATCTACAAGCAGAAGGGCGGAGGAGACCGGCAACTCAC
 AGACAGAGAGAAGGAGATCATCCGCCAGGCAGCGTGCAGCAGACCAAGGAGATGGACCTGAGCGTGGTG
 CGCCTCATGTTACAGCCTTCTCCCTGACAGCACTGGCAGCTTCACTCGGAGACTGGAGCCTGTGGTGT
 CAGACGCCATCTATGATAGCAAAGCCCCGAATGCATCCAACCTGAAAATCGTGAGAATGGACAGAACAGC
 AGGATGTGTGACGGGAGGGGAGGAGATTTACCTTCTGTGACAAGGTTTCAGAAAGATGACATCCAGATT
 CGGTTTTATGAAGAGGAAGAAAATGGCGGAGTTTGGGAAGGATTTGGGGACTTTTCCCCACAGATGTTT
 ATAGACAGTTTGCATTGTCTTCAAACGCCAAAGTATAAGGATGTCAACATTACAAAGCCAGCTTCCGT
 GTTTGTTTCAGCTTCGGAGGAAATCAGACCTGAAAAGTGAACCGAAACCTTTCTCTACTACCTGAA
 ATCAAAGACAAAGAGGAAGTCAAAGGAAACGCCAGAAGCTTATGCCGAACCTCTCGGACAGCTTCGGTG
 CGGGCAGTGGAGCGGGAGCCGGTGGTGGAGCATGTTTCGATAGTGGCGGTGGCGGAGGGATACCGGAAG
 CCTGGGCCAGGGTATGGCTACTCGAACTACGGATTTCCCTCCCTACGGTGGGATTACATTCATCCCGGA
 GTCACGAAATCCAACGCAGGGGTACCCATGGCACCATAAACACCAAATTTAAAAATGGCCCTAAAGATT
 GTGCCAAGAGTGTGACGAGGAGAGTCTGACTCTCCCTGAGAAGGAAACTGAAGGTGAAGGGCCAGCCT
 GCCCATGGCCTGCACCAAGAGCGGAACCCATCGCCTTGGCATCCACATGGAAGACAAGGAGCAGGACATG
 GGATTTACAGATAACCTCTTTCTTGAAGGCTCTGCAGCTCGCCAGGCGACACGCCAACGCCCTTTTCG
 ACTACGCAGTGACGGGGATGTGAAGATGTTGCTGGCCGTGCAACGCCATCTCACCGCCTGCAGGATGA
 GAATGGGGACAGTGTCTTACACTTAGCCATCATCCACCTCCACGCTCAGCTTGTGAGGGATCTGCTGGAA
 GTCACATCTGGTTGATCTCTGATGACATCATCAACATGAGAAATGACCTGTATCAGACACCTCTGCACT
 TGGCCGTGATACCAAGCAGGAAGATGTAGTAGAGGATTTGCTGAGGGTTGGGGCTGACCTGAGCCTTCT
 GGACCGCTGGGGCAACTCTGCTCCTGCACCTAGCTGCCAAAGAAGGACACGACAGAATCCTCAGCATCCTG
 CTCAAGAGCAGAAAAGCAGCGCCCTTATCGACCACCCCAATGGGGAAGGTCTAAATGCCATCCACATAG
 CTGTGATGAGCAATAGCCTGCCATGTCTGCTGCTGCTGGTGGCTGCCGGGCGAGAAGTCAATGCTCAGGA
 GCAGAAGTCTGGGCGCACAGCGCTGCACCTGGCCGTGGAGTACGACAACATCTCCTTGGCTGGCTGCCTG
 CTTCTGGAGGGTATGCCACGTGGACAGTACCACCTATGATGGGACTACACCTCTGCATATAGCGGCCG
 GAAGAGGGTCCACCAGACTGGCAGCTCTTCTCAAAGCAGCAGGAGCAGACCCCTGGTGGAGAACCTTGA
 GCCTCTCTATGACCTGGACGACTCTTGGGAGAAGGCTGGAGAAGATGAGGGAGTGGTCCAGGTACCACA
 CCCTTGACATGGCTGCCAACTGGCAGGATTTGACATACTAAATGGGAAACCGTATGAGCTGTGTTCA
 CATCTGATGATATACTACCACAAGGGGACATGAAGCAGCTGACAGAAGACACGAGGCTACAACCTGCAA
 ACTGCTGGAAATCCTGATCCAGACAAAACCTGGGCCACTCTGGCACAGAAGTTGGGTCTGGGGATACTG
 AACAAATGCCTTCCGGCTGAGTCTGCTCCTTCTAAAACCTCTCATGGACAACATGAGGTCTCTGGGGTA
 CCATCAAAGAGCTGATGGAGGCCCTGCAACAGATGGGCTACACAGAGGCCATTGAAGTATCCAGGCAGC
 CTTCCGCACCCCGCAACCACAGCCTCCAGCCCCGTGACCACTGCTCAGGTCCACTGTCTGCCTCTCTCG
 TCTTCTCCACGAGGCAGCACATAGATGAACTCCGGGATAGTGACAGCGTCTGTGACAGTGGTGTGGAGA
 CATCCTTCCGCAAACTCAGCTTTACAGAGTCTTACTGGAGACAGCCACTGCTATCTCTGAACAAAAT
 GCCCCACGGTTATGGGCAGGAAGGACCTATTGAAGGCAAAAT

ACGCGTACGCGGCCGCTCGAGCAGAAAACCTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA


Protein Sequence: >MR226792 protein sequence
 Red=Cloning site Green=Tags(s)

MADDDPYGTGQMFHLNTALTHSIFNAELYSPEIPLSTDGPYLQILEQPKQRGFRFRYVCEGPSHGGLPGA
 SSEKNKKSYPQVKICNYVGPQVIVQLVTNGKNIHLHAHSLVGKHCEDGVCTVTAGPKDMVYGFANLGIL
 HVTKKKVFETLEARMTEACIRGYNPGLLVHSDLAYLQAEGGDRQLTDREKEIIRQAAVQTKEMDLSV
 RLMFTAFLPDSTGFSFTRRLEPVVSDAIYDSKAPNASNLKIVRMDRTAGCVTGGEIYLLCDKVQKDDIQI
 RFYEEEEENGWVWEGGDFSPDVRQFAIVFKTPKYKDVNITKPASVQVLRKSDLETSEPKPFLYYPE
 IKDKKEEVQRKRQKLMNPFSDSFGGGSGAGAGGGMGFGSGGGGGSTGSPGPGYGSNYGFPYGGITFHFG
 VTKSNAGVTHGTINTKFKNGPKDCAKSDDEESLTLPEKETE GEGPSLPMACKTEPIALASTMEDKEQDM
 GFQDNLFLEKALQLARRHANALFDYAVTGDVKMLLAVQRHLTAVQDENGDSVLHLAIIHLHAQLVRDLLE
 VTSGLISDDIINMRNDLYQTPHLAVITKQEDVVEDLLRVGADLSLLDRWGNVSLHAAKEGHDRLSIL
 LKSRKAAPLIDHPNGEGLNAIHIIVMSNSLPCLLLLVAAGAEVNAQE QKSGRTALHLAVEYDNISLAGCL
 LLEGDAHVDSTTYDGTTPHLIAAGRGSTRLAALLKAAGADPLVENFEPLYDLDDSWEKAGEDEGVVPGTT
 PLDMAANWQVFDILNGKPYEPVFTSDDILPQGMKQLTEDTRLQLCKLLEIPDPDKNWATLAQKGLGIL
 NNAFRLSPAPSKTLMNDYEVSGGTIKELMEALQQMGYTEAIEVIQAAFRTATTASSPVTTAQVHCLPLS
 SSSTRQHIDELRSDSVCDSGVETSFRKLSFTESLTGDSPLL SLNKMPHYGYGQEGPIEGKI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:




ACCN: NM_008689
ORF Size: 2916 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_008689.2 , NP_032715.2
RefSeq Size:	4128 bp
RefSeq ORF:	2916 bp
Locus ID:	18033
UniProt ID:	P25799
Cytogenetics:	3 62.82 cM
MW:	105.6 kDa

Gene Summary:

NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. NF-kappa-B heterodimeric p65-p50 and RelB-p50 complexes are transcriptional activators. The NF-kappa-B p50-p50 homodimer is a transcriptional repressor, but can act as a transcriptional activator when associated with BCL3. NFKB1 appears to have dual functions such as cytoplasmic retention of attached NF-kappa-B proteins by p105 and generation of p50 by a cotranslational processing. The proteasome-mediated process ensures the production of both p50 and p105 and preserves their independent function, although processing of NFKB1/p105 also appears to occur post-translationally. p50 binds to the kappa-B consensus sequence 5'-GGRNNYYCC-3', located in the enhancer region of genes involved in immune response and acute phase reactions. Plays a role in the regulation of apoptosis. Isoform 5, isoform 6 and isoform 7 act as inhibitors of transactivation of p50 NF-kappa-B subunit, probably by sequestering it in the cytoplasm. Isoform 3 (p98) (but not p84 or p105) acts as a transactivator of NF-kappa-B-regulated gene expression. In a complex with MAP3K8, NFKB1/p105 represses MAP3K8-induced MAPK signaling; active MAP3K8 is released by proteasome-dependent degradation of NFKB1/p105.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR226792