

Product datasheet for MR226851

Aire (NM_009646) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Aire (NM_009646) Mouse Tagged ORF Clone

Tag: Myc-DDK

Symbol: Aire

Synonyms: MGC123374; MGC123375

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





ORF Nucleotide Sequence:

>MR226851 representing NM_009646
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ATGGCAGGTGGGGATGGAATGCTACGCCGTCTGCTGAGGCTGCACCGCACCGAGATCGCGGTGGCCATAG ACAGTGCCTTTCCGCTGCATGCTCTAGCCGACCACGACGTGGTCCCTGAGGACAAGTTCCAGGAGAC GCTCCGTCTGAAGGAGAAGGAAGGCTGCCCCCAGGCCTTCCACGCCCTGCTGTCCTGGCTCCTGACCCGG GACAGTGGGGCCATCCTGGATTTCTGGAGGATTCTCTTTAAGGACTACAATCTGGAGCGGTACAGCCGCC TGCATAGCATCCTGGACGGCTTCCCAAAAGATGTGGACCTAAACCAGTCCCGGAAAGGGAGAAAGCCCCT TGCTGGTCCCAAGGCCGCGGTACTGCCACCCAGACCCCCACCAAGAGAAAAGCACTGGAGGAGCCTCGA GCCACCCACCAGCAACTCTGGCCTCAAAGAGCGTCTCCAGCCCAGGCTCCCACCTGAAGACTAAGCCCC CTAAGAAGCCAGATGGCAACTTGGAGTCACAGCACCTTCCTCTTGGAAACGGAATTCAGACCATGGCAGC TTCTGTCCAGAGAGCTGTGACCGTGGCCTCTGGGGATGTTCCAGGAACCCGAGGGGCCGTGGAAGGGATC CTTATCCAGCAGGTGTTTGAGTCAGGAAGATCCAAGAAGTGCATTCAGGTTGGGGGAGAGTTTTATACAC CCAACAAGTTCGAAGACCCCAGTGGCAATTTGAAGAACAAGGCCCGGAGTGGTAGCAGCCTAAAGCCAGT GGTCCGAGCCAAGGGAGCCCAGGTCACTATACCTGGTAGAGATGAGCAGAAAGTGGGCCAGCAGTGTGGG GTTCCTCCCCTTCCATCCCCCAGTGAGCCCCAGGTTAACCAGAACGAGGATGAGTGTGCCGTGTGCC ACGACGGAGGTGAGCTCATCTGTTGTGACGGCTGTCCCCGGGCCTTCCACCTGGCTTGCCTGTCCCCACC TCTGCAGGAGATCCCCAGTGGCCTCTGGAGATGCTCCTGCTGCCTCCAGGGCAGAGTCCAACAGAACCTG TCCCAGCCTGAGGTGTCCAGGCCCCCGGAGCTACCTGCAGAGACCCCGATCCTCGTGGGACTGAGGTCAG CTTCAGAGAAAACCAGGGGCCCATCCAGGGAGCTCAAAGCCAGCTCTGATGCTGCTGTCACATATGTGAA CCTGCTGGCCCCGCACCCTGCAGCTCCTCTGCTGGAGCCTTCAGCACTGTGCCCTCTACTGAGTGCTGGG CAATCTCCGCTGCAAATCCTGCTCTGCAGACTCGACTCCCACGCCAGGCACACCGGGCGAAGCTGTACCC ACCTCTGGGCCCCGTCCAGCACCTGGGCTTGCCAAGGTAGGGGACGACTCTGCTAGTCACGACCCTGTTC TACATAGGGACGACCTGGAGTCCCTCCTCAATGAGCACTCATTTGACGGCATCCTGCAGTGGGCCATCCA GAGCATGTCACGCCCGCTGGCCGAGACACCACCCTTCTCTCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:

>MR226851 representing NM_009646 Red=Cloning site Green=Tags(s)

MAGGDGMLRRLLRLHRTEIAVAIDSAFPLLHALADHDVVPEDKFQETLRLKEKEGCPQAFHALLSWLLTR DSGAILDFWRILFKDYNLERYSRLHSILDGFPKDVDLNQSRKGRKPLAGPKAAVLPPRPPTKRKALEEPR ATPPATLASKSVSSPGSHLKTKPPKKPDGNLESQHLPLGNGIQTMAASVQRAVTVASGDVPGTRGAVEGI LIQQVFESGRSKKCIQVGGEFYTPNKFEDPSGNLKNKARSGSSLKPVVRAKGAQVTIPGRDEQKVGQQCG VPPLPSLPSEPQVNQNEDECAVCHDGGELICCDGCPRAFHLACLSPPLQEIPSGLWRCSCCLQGRVQQNL SQPEVSRPPELPAETPILVGLRSASEKTRGPSRELKASSDAAVTYVNLLAPHPAAPLLEPSALCPLLSAG NEGRPGPAPSARCSVCGDGTEVLRCAHCAAAFHWRCHFPTAAARPGTNLRCKSCSADSTPTPGTPGEAVP TSGPRPAPGLAKVGDDSASHDPVLHRDDLESLLNEHSFDGILQWAIQSMSRPLAETPPFSS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Restriction Sites:

Sgfl-Mlul



Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_009646

ORF Size: 1653 bp

OTI Disclaimer:

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

OTI Annotation:

This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.



RefSeq: <u>NM 009646.1, NM 009646.2, NP 033776.1</u>

 RefSeq Size:
 1993 bp

 RefSeq ORF:
 1659 bp

 Locus ID:
 11634

 UniProt ID:
 Q9Z0E3


Cytogenetics: 10 39.72 cM **MW:** 59.5 kDa

Gene Summary:

Transcription factor playing an essential role to promote self-tolerance in the thymus by regulating the expression of a wide array of self-antigens that have the commonality of being tissue-restricted in their expression pattern in the periphery, called tissue restricted antigens (TRA) (Probable). Binds to G-doublets in an A/T-rich environment; the preferred motif is a tandem repeat of 5'-. ATTGGTTA-3' combined with a 5'-TTATTA-3' box. Binds to nucleosomes (By similarity). Binds to chromatin and interacts selectively with histone H3 that is not methylated at 'Lys-4', not phosphorylated at 'Thr-3' and not methylated at 'Arg-2'. Functions as a sensor of histone H3 modifications that are important for the epigenetic regulation of gene expression. Mainly expressed by medullary thymic epithelial cells (mTECs), induces the expression of thousands of tissue-restricted proteins, which are presented on major histocompatibility complex class I (MHC-I) and MHC-II molecules to developing T-cells percolating through the thymic medulla (By similarity). Also induces self-tolerance through other mechanisms such as the regulation of the mTEC differentiation program (PubMed:19015306). Controls the medullary accumulation of thymic dendritic cells and the development of regulatory T-cell through the regulation of XCL1 expression (PubMed:21300913). Regulates the production of CCR4 and CCR7 ligands in medullary thymic epithelial cells and alters the coordinated maturation and migration of thymocytes (PubMed:19923453). In thimic B-cells, allows the presentation of licensing-dependent endogenous self-anitgen for negative selection (PubMed:26070482). In secondary lymphoid organs, induces functional inactivation of CD4(+) T-cells. Expressed by a distinct bone marrow-derived population, induces self-tolerance through a mechanism that does not require regulatory T-cells and is resitant to innate inflammatory stimuli (PubMed:23993652). [UniProtKB/Swiss-Prot Function]



Product images:



Circular map for MR226851