

Product datasheet for **MR226922**

Prkcq (NM_008859) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Prkcq (NM_008859) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Prkcq
Synonyms:	A130035A12Rik; AW494342; PKC-0; PKC-theta; Pkcq; PKCtheta
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>MR226922 representing NM_008859
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGTCACCGTTTCTCGAATCGGTTTATCCAACCTTGACTGTGGGACCTGCCAAGCTTGTACGGGAGAGG
 CAGTGAACCCCTACTGCGCTGTGCTTGTCAAAGAGTATGTGGAATCAGAAAATGGGCAGATGTACATCCA
 GAAAAAGCCAACCATGTACCCACCTTGGGACAGCACCTTTGACGCCACATTAACAAGGGAAGGGTGATG
 CAGATCATCGTGAAGGGCAAGAATGTAGACCTCATCTCAGAAAACAACCGTGAACCTACTCCCTGGCGG
 AGAGATGCCGCAAGAACAATGGGCGGACAGAAATATGGTTAGAGCTGAAACCTCAAGGCCGAATGCTAAT
 GAATGCAAGATACTTTCTGAAAATGAGTGACACAAAGGACATGAGTGAGTTTGAAGTGAAGGATCTTT
 GCACTGCATCAGCGCCGAGGAGCCATCAAACAGGCCAAAGTCCACCATGTCAAGTGCACGATTCACGG
 CCACCTTTTTCCCTCAACCCACATTTTGTCTGTCTGCCATGAATTTGTCTGGGGCTGAACAAGCAGGG
 TTACCAGTGCCGACAGTGAATGCAGCGATTACAAGAAGTGCATTGATAAAGTGTAGCCAAGTGACACA
 GGATCCGCAATCAATAGCCGAGAAAACCATGTTCCATAAGGAGAGATTCAAGATCGACATGCCACACAGAT
 TCAAAGTCTACAACACAAAGAGTCCAACCTTCTGTGAGCACTGTGGTACCTGCTCTGGGGCTGGCGAG
 GCAAGGACTCAAATGTGATGCATGTGGCATGAACGTCCACCACCGATGCCAGACAAAAGTTGCCAATCTT
 TGTGGTATAAACCAGAAGCTAATGGCTGAAGCACTAGCGATGATTGAAAGCACCAACAGGCTCGCTCCT
 TACGAGATTCAGAACACATCTCCGAGAAGGCCAGTTGAAATTTGGTCTCCCATGCTCCACCAAAAAACGA
 AACCCAGGCCACCATGCGTACCAACACCTGGGAAAAGAGAACCCAGGGCATTTCCTGGGATTCCTTTTG
 GATGGGTCAAATAAATCGGCCGGTCTCCTGAACCCGAAGTGAAGTGCATGCGCAGGACTTACTGCAGCTGA
 AACTGAAGATCGATGACTTCACTCCTGCACAAGATGTTGGGAAAAGGAAGTTTGGCAAGGCTTCCTGGC
 AGAGTTCAAGAGAACCAATCAGTTTTTCGCAATAAAAGCCTTAAAGAAAGATGTGGTGTGATGGATGAT
 GACGTCGAGTGTACAATGGTGGAAAAGAGGGTTCTGTCTTGGCATGGGAGCATCCATTTCTAACACACA
 GTTTCTGCACATTCAGACCAAGGAAAATCTCTTTTTCTGTGATGGAGTATCTCAATGGAGGCGACTTAAT
 GTACCACATCAAAGTTGCCACAAATTTGATCTTCCAGAGCCACGTTTTATGCTGCTGAGGTCATCCTT
 GGTCTGCAGTTCCTTCATTCAAAGGAATTGTCTACAGGGACCTGAAGCTTGATAATATCCTGTTAGACA
 GAGATGGACATATCAAAATAGCAGACTTTGGGATGTGCAAAGAGAACATGCTAGGAGATGCGAAGACAAA
 TACTTTCTGTGGAACCTCTGACTACATTGCTCCGGAGATCTTGTGGTCAAGTACAACCATTCGCTC
 GACTGGTGGTCTTCGGGGTCTCGTTTATGAGATGCTGATTGGCCAGTCCCCCTCCACGGGCAGGACG
 AGGAGGAGCTGTCCACTCCATCCGCATGGACAACCCCTTCTACCCGAGGTGGCTCGAAAAGGAGGCCAA
 GGACCTTCTAGTGAAGCTTTTTGTGAGAGAACCTGAGAAGAGGCTGGGAGTGAGAGGAGACATCCGCCAG
 CATCCTTTGTTTCGAGAGATCAACTGGGAAGAGCTTGAAGGAAAAGAGATTGACCCACCTTCAGACCAA
 AAGTGAATCACCATATGACTGTAGCAATTTGACAAGGAATTCCTAAGTGAAGAACCCCGGCTATCATT
 CGCCGACAGACTCATCAACAGCATGGACCAGAACATGTTTCAGCAACTTTTCTTCATTAACCCAGGG
 ATGGAGACTCTCATTGCTCC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA


Protein Sequence: >MR226922 representing NM_008859
 Red=Cloning site Green=Tags(s)

```
MSPFLRIGLSNFDCGTCQACQGEAVNPYCAVLVKEYVESENGQMYIQKKPTMYPPWDSTFDAHINKGRVM
QIIYKGNVDLISETTVELYSLAERCNRNNGRTEIWLKLPQGRMLMNARYFLEMSDTKDMSEFENEFF
ALHQRRAIKQAKVHVHKCHEFTATFFPQPTFCSVCHEFVWGLNKQGYQCRQCNAAIHKKIDKVIKCT
GSAINSRETMFHKERFKIDMPHRFKVYNYKSPTFCEHCGTLLWGLARQGLKCDACGMNVVHRCQTKVANL
CGINQKLMAEALAMIESTQQARSLRDSEHIFREGPVEIGLPCSTKNETRPPCVPTPGKREPOGISWDSPL
DGSNKSAGPPEPEVSMRRTSLQLKLIKIDDFILHKMLGKGSFGKVFLEAFKRTNQFFAIKALKKDVVLMDD
DVECTMVEKRVLSLAWHEPFLTHMFCFQTKENLFFVMEYLNNGDLMYHIQSCHKFDLSRATFYAAEVIL
GLQFLHSKGIYVRDLKLDNILLDRDGHKIDAFGMCKENMLGDAKTNTFCGTPDYIAPEILLGQKYNHSV
DWSFGVLVYEMLIGQSPFHGQDEEELFHSIRMDNPFYPRWLERAKDLLVKLFVREPEKRLGVRGDIRQ
HPLFREINWEELERKEIDPPFRPKVKSPYDCSNFDKEFLSEKPRLSFADRALINSMDQNMFSNFSFINPG
METLICs
```

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_008859

ORF Size: 2121 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_008859.2](#), [NP_032885.1](#)

RefSeq Size: 3313 bp

RefSeq ORF: 2124 bp

Locus ID: 18761

UniProt ID: [Q02111](#)


Cytogenetics: 2 8.42 cM

MW: 82 kDa

Gene Summary:

Calcium-independent, phospholipid- and diacylglycerol (DAG)-dependent serine/threonine-protein kinase that mediates non-redundant functions in T-cell receptor (TCR) signaling, including T-cells activation, proliferation, differentiation and survival, by mediating activation of multiple transcription factors such as NF-kappa-B, JUN, NFATC1 and NFATC2. In TCR-CD3/CD28-co-stimulated T-cells, is required for the activation of NF-kappa-B and JUN, which in turn are essential for IL2 production, and participates in the calcium-dependent NFATC1 and NFATC2 transactivation. Mediates the activation of the canonical NF-kappa-B pathway (NFKB1) by direct phosphorylation of CARD11 on several serine residues, inducing CARD11 association with lipid rafts and recruitment of the BCL10-MALT1 complex, which then activates IKK complex, resulting in nuclear translocation and activation of NFKB1. May also play an indirect role in activation of the non-canonical NF-kappa-B (NFKB2) pathway. In the signaling pathway leading to JUN activation, acts by phosphorylating the mediator STK39/SPAK and may not act through MAP kinases signaling. Plays a critical role in TCR/CD28-induced NFATC1 and NFATC2 transactivation by participating in the regulation of reduced inositol 1,4,5-trisphosphate generation and intracellular calcium mobilization. After costimulation of T-cells through CD28 can phosphorylate CBLB and is required for the ubiquitination and subsequent degradation of CBLB, which is a prerequisite for the activation of TCR. During T-cells differentiation, plays an important role in the development of T-helper 2 (Th2) cells following immune and inflammatory responses, and, in the development of inflammatory autoimmune diseases, is necessary for the activation of IL17-producing Th17 cells. May play a minor role in Th1 response. Upon TCR stimulation, mediates T-cell protective survival signal by phosphorylating BAD, thus protecting T-cells from BAD-induced apoptosis, and by up-regulating BCL-X(L)/BCL2L1 levels through NF-kappa-B and JUN pathways. In platelets, regulates signal transduction downstream of the ITGA2B, CD36/GP4, F2R/PAR1 and F2RL3/PAR4 receptors, playing a positive role in 'outside-in' signaling and granule secretion signal transduction. May relay signals from the activated ITGA2B receptor by regulating the uncoupling of WASP and WIPF1, thereby permitting the regulation of actin filament nucleation and branching activity of the Arp2/3 complex. May mediate inhibitory effects of free fatty acids on insulin signaling by phosphorylating IRS1, which in turn blocks IRS1 tyrosine phosphorylation and downstream activation of the PI3K/AKT pathway. Phosphorylates MSN (moesin) in the presence of phosphatidylglycerol or phosphatidylinositol. Phosphorylates PDPK1 at 'Ser-504' and 'Ser-532' and negatively regulates its ability to phosphorylate PKB/AKT1.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR226922