

Product datasheet for MR226996L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Nf2 (NM_010898) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Nf2 (NM_010898) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Nf2

Synonyms: merlin

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_010898

ORF Size: 1788 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR226996).

•

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 010898.3, NP 035028.2

 RefSeq Size:
 4708 bp

 RefSeq ORF:
 1791 bp

 Locus ID:
 18016

 UniProt ID:
 P46662

Cytogenetics: 11 3.0 cM







Gene Summary:

Probable regulator of the Hippo/SWH (Sav/Wts/Hpo) signaling pathway, a signaling pathway that plays a pivotal role in tumor suppression by restricting proliferation and promoting apoptosis. Along with WWC1 can synergistically induce the phosphorylation of LATS1 and LATS2 and can probably function in the regulation of the Hippo/SWH (Sav/Wts/Hpo) signaling pathway. May act as a membrane stabilizing protein. May inhibit PI3 kinase by binding to AGAP2 and impairing its stimulating activity. Suppresses cell proliferation and tumorigenesis by inhibiting the CUL4A-RBX1-DDB1-VprBP/DCAF1 E3 ubiquitin-protein ligase complex (By similarity). Plays a role in lens development and is required for complete fiber cell terminal differentiation, maintenance of cell polarity and separation of the lens vesicle from the corneal epithelium.[UniProtKB/Swiss-Prot Function]