

## Product datasheet for MR227021L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Txnip (NM\_001009935) Mouse Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Txnip (NM\_001009935) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Txnip

**Synonyms:** 1200008J08Rik; AA682105; Hyplip1; Tbp-2; THIF; VDUP1

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001009935

ORF Size: 1191 bp

**ORF Nucleotide** 

\_. . . . . .

Sequence:

The ORF insert of this clone is exactly the same as(MR227021).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001009935.2

 RefSeq Size:
 2801 bp

 RefSeq ORF:
 1194 bp

 Locus ID:
 56338

 UniProt ID:
 Q8BG60

Cytogenetics: 3 41.93 cM





## **Gene Summary:**

May act as an oxidative stress mediator by inhibiting thioredoxin activity or by limiting its bioavailability. Interacts with COPS5 and restores COPS5-induced suppression of CDKN1B stability, blocking the COPS5-mediated translocation of CDKN1B from the nucleus to the cytoplasm. Inhibits the proteasomal degradation of DDIT4, and thereby contributes to the inhibition of the mammalian target of rapamycin complex 1 (mTORC1) (By similarity). Functions as a transcriptional repressor, possibly by acting as a bridge molecule between transcription factors and corepressor complexes, and over-expression will induce G0/G1 cell cycle arrest. Required for the maturation of natural killer cells. Acts as a suppressor of tumor cell growth.[UniProtKB/Swiss-Prot Function]