

Product datasheet for **MR227044L4V**

Pecam1 (NM_008816) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Pecam1 (NM_008816) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Pecam1
Synonyms:	C85791; Cd31; Pecam; PECAM-1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_008816
ORF Size:	2148 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR227044).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_008816.2
RefSeq Size:	3305 bp
RefSeq ORF:	2151 bp
Locus ID:	18613
UniProt ID:	Q08481
Cytogenetics:	11 E1



[View online »](#)

Gene Summary:

Cell adhesion molecule which is required for leukocyte transendothelial migration (TEM) under most inflammatory conditions (By similarity). Tyr-679 plays a critical role in TEM and is required for efficient trafficking of PECAM1 to and from the lateral border recycling compartment (LBRC) and is also essential for the LBRC membrane to be targeted around migrating leukocytes (By similarity). Trans-homophilic interaction may play a role in endothelial cell-cell adhesion via cell junctions (By similarity). Heterophilic interaction with CD177 plays a role in transendothelial migration of neutrophils (By similarity). Homophilic ligation of PECAM1 prevents macrophage-mediated phagocytosis of neighboring viable leukocytes by transmitting a detachment signal (By similarity). Promotes macrophage-mediated phagocytosis of apoptotic leukocytes by tethering them to the phagocytic cells; PECAM1-mediated detachment signal appears to be disabled in apoptotic leukocytes (By similarity). Modulates bradykinin receptor BDKRB2 activation (By similarity). Regulates bradykinin- and hyperosmotic shock-induced ERK1/2 activation in endothelial cells (By similarity). Induces susceptibility to atherosclerosis (PubMed:19048083).[UniProtKB/Swiss-Prot Function]