

Product datasheet for MR227135L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Grin2a (NM_008170) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Grin2a (NM_008170) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Grin2a

Synonyms: GluN2A; NMDAR2A; NR2A

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_008170 **ORF Size:** 4392 bp

ORF Nucleotide

OTI Disclaimer:

- - - - 1

Sequence:

The ORF insert of this clone is exactly the same as(MR227135).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 008170.2

 RefSeq Size:
 4512 bp

 RefSeq ORF:
 4395 bp

 Locus ID:
 14811

 UniProt ID:
 P35436

Cytogenetics: 16 5.28 cM





Gene Summary:

Component of NMDA receptor complexes that function as heterotetrameric, ligand-gated ion channels with high calcium permeability and voltage-dependent sensitivity to magnesium (PubMed:1374164). Channel activation requires binding of the neurotransmitter glutamate to the epsilon subunit, glycine binding to the zeta subunit, plus membrane depolarization to eliminate channel inhibition by Mg(2+). Sensitivity to glutamate and channel kinetics depend on the subunit composition; channels containing GRIN1 and GRIN2A have higher sensitivity to glutamate and faster kinetics than channels formed by GRIN1 and GRIN2B (By similarity). Contributes to the slow phase of excitatory postsynaptic current, long-term synaptic potentiation, and learning (PubMed:7816096, PubMed:8987814).[UniProtKB/Swiss-Prot Function]