

Product datasheet for **MR227160L3V**

Trpv1 (NM_001001445) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Trpv1 (NM_001001445) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Trpv1
Synonyms:	OTRPC1; TRPV1alpha; TRPV1beta; VR-1; Vr1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001001445
ORF Size:	2517 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR227160).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_001001445.1
RefSeq Size:	2520 bp
RefSeq ORF:	2520 bp
Locus ID:	193034
UniProt ID:	Q704Y3
Cytogenetics:	11 45.25 cM



[View online »](#)

Gene Summary:

Ligand-activated non-selective calcium permeant cation channel involved in detection of noxious chemical and thermal stimuli (PubMed:15194687, PubMed:15489017). Seems to mediate proton influx and may be involved in intracellular acidosis in nociceptive neurons. Involved in mediation of inflammatory pain and hyperalgesia (PubMed:10764638). Sensitized by a phosphatidylinositol second messenger system activated by receptor tyrosine kinases, which involves PKC isozymes and PCL. Activation by vanilloids, like capsaicin, and temperatures higher than 42 degrees Celsius, exhibits a time- and Ca(2+)-dependent outward rectification, followed by a long-lasting refractory state. Mild extracellular acidic pH (6.5) potentiates channel activation by noxious heat and vanilloids, whereas acidic conditions (pH <6) directly activate the channel. Can be activated by endogenous compounds, including 12-hydroperoxytetraenoic acid and bradykinin. Acts as ionotropic endocannabinoid receptor with central neuromodulatory effects. Triggers a form of long-term depression (TRPV1-LTD) mediated by the endocannabinoid anandamine in the hippocampus and nucleus accumbens by affecting AMPA receptors endocytosis (By similarity).[UniProtKB/Swiss-Prot Function]