

## Product datasheet for **MR227181**

### **Irak1 (NM\_001177975) Mouse Tagged ORF Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	Irak1 (NM_001177975) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Irak1
Synonyms:	AA408924; Il1rak; IRAK; IRAK-1; IRAK1-S; IRAK1b; mPLK; Plpk
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**ORF Nucleotide Sequence:**

>MR227181 representing NM\_001177975  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGGATCGCC**

ATGGCCGGGGGGCCGGGCCCGGGGAGCCTGTGGTTCCAGGCGCCAGCACTTCTTGTACGAGGTGCCAC  
 CCTGGGTTATGTGCCGCTTCTACAAAGTGATGGACGCCCTGGAGCCCGCGACTGGTGCCAGTTCGCGGC  
 CTTGATCGTGCGCGACCAGACAGAGCTGCGGCTGTGCGAGCGCTCCGAGCAGCGCACAGCCAGTGTCTG  
 TGGCCCTGGATCAACCGCAACGCGCGCGTAGTGACCTCGTTCACATCCTCACGCACCTGCAGCTGCTGC  
 GTGCGAGGGACATCATCACAGCCTGGCACCCCTCCTGCCCCGTTGTGCCCAAGCACCGCTGCCCAAG  
 GCCCAGCAGCATCTGCGAGGCTCTGAGGCCGGGACTGGAGCCCCGGAAATTGCAGTCTCTGCCTCC  
 ACCTTCTCTCCCGAGCTTTCCAGGCTCCAGACCCATTCTGAGTCAGAGCTCCTCCAGGTTCCACTCC  
 CTGTTTCCCTCGGGCCACCCTACCCTTTCAGCCCTTCTCCACCAAGCAGTCAAGCCCAGAGAGCCC  
 AGTGTCTGGCCTCAAAGAGCCCATCCCTCCCGTTTTGTGGCCCTTCTGTGAGATTTCCAAGGCACC  
 TGCAACTTCTCTGAAGAACTCAGGATTGGAGAGGGTGGTTTTGGATGTGTGTACCGAGCAGTTCATGAGAA  
 ATACTACATATGCTGTGAAGAGACTGAAGGAGGAGCTGACCTAGAGTGGACTATGGTGAACAGAGCTT  
 CTTAACAGAGGTGGAACAGCTATCAAGGTTTCTGACCCAAATATCGTAGACTTTGCTGGCTACTGTGCA  
 GAGAGTGGCTTACTGCCTTGTATGGCTTCTTGCCCAATGGCTCCTTGAGGATCAGCTCCACCTTC  
 AGACCCAGGCCTGCTCCCACTTCTGGCCTCAACGACTGGACATTCCTTGGGCACAGCCCGGCTAT  
 TCAGTTTTTACATCAGGATAGCCCCAGCCTTATCCATGGAGACATCAAGAGTTCTAACGTGCTTCTGGAT  
 GAGAGACTGATGCCAAGCTGGGAGACTTTGGCCTGGCTCGTTTTAGCCGCTTTGCGGGGGCCAAAGCAA  
 GCCAGAGCAGTACTGTGGCCCGGACTTCCACAGTTCGAGGTACCCTCGCCTACTTGCCTGAGGAGTACAT  
 CAAGACAGGCCGACTGGCTGTGGACACCGATACCTTCAGCTTTGGGGTGGTAATACTGGAGACCCTTGCT  
 GGTGAGAGGGCTGTGAGGACACAAGGTGCAAAGACCAAGTATTTGAAAGACCTGATTGAAGATGAGGCTG  
 AAGAGGCTGGAGTGACCCTGAAAAGCACCCAGCCTACTCTGTGGGTGGGTGTAGCCACGGATGCTTGGGC  
 TGCTCCAATTGCTGCCAGATCTATAAGAAGCACCTGGACTCCAGACCTGGGCCCTGCCACCCAGTTG  
 GGCCTGGCCCTGGCTCAACTAGCTTGTGCTGCATGCACCGTGGGCCAAGAAGAGGCCCCCATGACCC  
 AGGTATAAAGAGACTAGAAGGGCTTCAGGCAGGGCCTCCCTGGGAGCTAGAGGTTGCCGGCCATGGCTC  
 CCCTTCCCACAGGAGAACTCCTACATGTCTACCACTGGCAGTGGCCAGAGTGGGGATGAACCATGGCAG  
 CCTCTAGTAGTGACCACAAGAGCCCAGCCAGGCTGCCAGCAACTCCAGAGAAGTCCCAACCAGCCAG  
 TGGAAAGTGATGAGAGTGTCCCGGCCTCTCTGCTACCCTGCATTCTGGCACTTGACTCCAGGTTCCCA  
 CCCAAGGCCAGCGTCTTCAGAGAGGCTAGCTGTACCCAGGGAGGCACTACCAGAGAATCAAGTGTGAGG  
 AGTAGCCAGGCTTCCAGCCTACAACCATGGAAGGCTCACCCACGGGCAGCTCATCCCTGCTGTCATCAG  
 AACCACCACAGATCATCAACCCAGCCCAGCAGAAGATGGTACAAAAGCTGGCTCTTTATGAAGAAGG  
 GGTCTTGATAGCCTGCAACTGCTGTATCAGGCTTTTTCCAGGCTTGGATTTAGAACCTGAAAAGAGC  
 CAGGGACCTGAAGAAAGTGTGAATTCAGAGC


**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA



<b>OTI Disclaimer:</b>	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:custsupport@origene.com">custsupport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a></p>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<u>NM_001177975.1, NP_001171446.1</u>
<b>RefSeq Size:</b>	3835 bp
<b>RefSeq ORF:</b>	2136 bp
<b>Locus ID:</b>	16179
<b>Cytogenetics:</b>	X 37.61 cM
<b>MW:</b>	77.4 kDa

**Gene Summary:**

Serine/threonine-protein kinase that plays a critical role in initiating innate immune response against foreign pathogens. Involved in Toll-like receptor (TLR) and IL-1R signaling pathways. Is rapidly recruited by MYD88 to the receptor-signaling complex upon TLR activation. Association with MYD88 leads to IRAK1 phosphorylation by IRAK4 and subsequent autophosphorylation and kinase activation. Phosphorylates E3 ubiquitin ligases Pellino proteins (PELI1, PELI2 and PELI3) to promote pellino-mediated polyubiquitination of IRAK1. Then, the ubiquitin-binding domain of IKBKG/NEMO binds to polyubiquitinated IRAK1 bringing together the IRAK1-MAP3K7/TAK1-TRAF6 complex and the NEMO-IKKA-IKKB complex. In turn, MAP3K7/TAK1 activates IKKs (CHUK/IKKA and IKBKB/IKKB) leading to NF-kappa-B nuclear translocation and activation. Alternatively, phosphorylates TIRAP to promote its ubiquitination and subsequent degradation. Phosphorylates the interferon regulatory factor 7 (IRF7) to induce its activation and translocation to the nucleus, resulting in transcriptional activation of type I IFN genes, which drive the cell in an antiviral state. When sumoylated, translocates to the nucleus and phosphorylates STAT3 (By similarity). [UniProtKB/Swiss-Prot Function]

**Product images:**

Circular map for MR227181