

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for MR227188L3V

S100b (NM_009115) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

| Product Type: | Lentiviral Particles |
|------------------------------|---|
| Product Name: | S100b (NM_009115) Mouse Tagged ORF Clone Lentiviral Particle |
| Symbol: | S100b |
| Synonyms: | Al850290; Bpb |
| Mammalian Cell Selection: | Puromycin |
| Vector: | pLenti-C-Myc-DDK-P2A-Puro (PS100092) |
| Tag: | Myc-DDK |
| ACCN: | NM_009115 |
| ORF Size: | 279 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(MR227188). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u> |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | <u>NM 009115.3, NP 033141.1</u> |
| RefSeq Size: | 1676 bp |
| RefSeq ORF: | 279 bp |
| Locus ID: | 20203 |
| UniProt ID: | <u>P50114</u> |
| Cytogenetics: | 10 38.76 cM |
| | |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

S100b (NM_009115) Mouse Tagged ORF Clone Lentiviral Particle – MR227188L3V

Gene Summary:Weakly binds calcium but binds zinc very tightly-distinct binding sites with different affinities
exist for both ions on each monomer. Physiological concentrations of potassium ion
antagonize the binding of both divalent cations, especially affecting high-affinity calcium-
binding sites. Binds to and initiates the activation of STK38 by releasing autoinhibitory
intramolecular interactions within the kinase. Interaction with AGER after myocardial
infarction may play a role in myocyte apoptosis by activating ERK1/2 and p53/TP53 signaling.
Could assist ATAD3A cytoplasmic processing, preventing aggregation and favoring
mitochondrial localization. May mediate calcium-dependent regulation on many physiological
processes by interacting with other proteins, such as TPR-containing proteins, and
modulating their activity (By similarity).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US