

Product datasheet for MR227208

Rag1 (NM_009019) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Rag1 (NM_009019) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Rag1
Synonyms:	Rag-1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>MR227208 representing NM_009019 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ATGGCTGCCTCCTTGCCGTCTACCCTGAGCTTCAGTTCTGCACCCGATGAAATCAACACCCACAAATCA
AATTTTCCGAGTGGAAATTAAGCTGTTTAGGGTGAATCCTTTGAAAAGGCACCCGAAGAAGCACAGAA
GGAGAAGGATTCCTCAGAGGGGAAACCTTACCTAGAACAGTCTCCAGTAGTTCAGAGAAGCCTGGTGGT
CAGAACTCAATTCTGACTCAACGAGCACTGAAACTCCATCCTAAATTTTCAAAGAAATTCATGCTGATG
GGAAGTCAAGCGACAAAGCAGTTCACCAAGCCAGGCTTAGACACTTCTGCCGCATCTGTGGGAATCGTTT
CAAGAGTGACGGGCACAGCCGGAGATACCCAGTCCACGGGCCCGTGGACGCTAAAACCCAAAGTCTTTTC
CGAAAGAAGGAAAAAGAGTCACTTCTGGCCAGACCTCATTGCCAGGATTTTCCGGATCGACGTGAAGG
CAGATGTTGACTCCATCCACCCGACGGAATTCTGCCATGACTGTTGGAGCATCATGCACAGAAAAGTTCAG
CAGTTCACACAGTCAAGTCTACTTCCCAAGGAAAGTGACCGTGGAGTGGCACCCCCACACACCCGCTCTGT
GACATCTGTTTTACTGCCATCGGGGACTCAAGAGGAAGAGACATCAGCCCAATGTGCAGCTCAGCAAGA
AACTAAAACTGTGCTCAACCACGCGAGACGGGACCGTCGCAAGAGAAGTCAAGGCTAGGGTCAAGCAAG
GGAAGTCTGAAGAAGATCTCCAAGTGCAGTAAAGTTCATCTCAGTACCAAGCTTCTTGGCGTGGACTTC
CCAGCACACTTTGTGAAATCCATCTCCTGCCAGATATGCGAACACATTCTGGCTGATCCCGTGGAGACCA
GCTGCAAGCATCTATTCTGTAGGATCTGCATTCTCAGATGTCTCAAAGTCATGGGCAGCTATTGTCCCTC
TTGCCGATATCCGTGCTTCCCTACTGACCTGGAGAGCCAGTGAAGTCCCTTCTGAACATCTTGAATTCT
CTCATGGTCAAGTGTCCCGCGCAAGATTGCAATGAGGAAGTGAAGTCTGGAAAAATAAACCACCATGTGT
CAAGCCACAAAGAATCTAAAGAGACTTTGGTGCAATCAATAAAGGGGGACGGCCCGCCAGCATCTCCT
GTCAGTACGAGAAGGGCGCAGAAACATCGGCTGAGGGAGCTCAAGATCAAGTCAAAGAATTTGCTGAC
AAAGAAGAAGGTGGAGATGTGAAGGCTGTCTGCTTGACATTGTTTCTCCTGGCACTGAGGGCGAGGAATG
AGCACAGGCAAGCTGATGAATTAGAGGCCATCATGCAAGGCAGGGGCTCCGGGCTTCAACCAGCTGTTTG
CTTGCCATCCGTGCAATACCTTCTCAGCTGTAGCCAATACCATAAGATGTACAGGACTGTGAAAGCT



[View online >](#)

ATCACTGGGAGGCAGATTTTTCAACCTTTGCATGCTCTTCGGAATGCCGAGAAAGTCCTTCTGCCAGGCT
 ACCATCCCTTTGAGTGGCAGCCCCACTGAAGAATGTGTCTCCAGAAGTATGTTGGAATTATTGATGG
 GCTGTCTGGACTTGCCTCCTCTGTGGATGAGTACCCAGTAGATACCATTGCGAAGAGGTTCCGCTACGAC
 TCTGCTTTGGTGTCTGCTTTGATGGACATGGAAGAAGACATCTTGGAAGGCATGAGATCCCAAGATCTTG
 ATGACTACCTGAATGGTCCCTTACAGTGGTGGTAAAGGAGTCTTGCATGGAATGGGGGATGTGAGTGA
 GAAGCACGGGAGTGGGCCCGAGTTCAGAAAAAGGCCGTTCTGTTCTCTTTCACAGTATGAGAATTACG
 ATAGAGCATGGTTCACAGAACGTGAAGGTGTTTGGGAACCCAAGCCCAATTCTGAAGTGTGTTGCAAGC
 CGTTGTGCTTATGCTGGCAGATGAGTCTGACCATGAGACCCCTTACTGCTATTCTAAGCCCCCTCATTGC
 CGAGAGGGAGGCCATGAAGAGCAGTGAATTAACGCTGGAGATGGGAGGCATCCCCAGGACTTTTAAATTC
 ATCTTCAGGGGCACTGGATACGATGAAAAACTTGTCCGGGAAGTAGAAGGCTTGAAGCTTCTGGCTCAG
 TCTACACTGTACTCTGTGACACCACCCGTTTGAAGCCTCTCAGAATCTTGTCTTCCACTCCATAAC
 CAGAAGCCACGCCGAGAACCTGCAGCGTATGAGGTCTGGCGGTCCAATCCGTATCATGAGTCCGTGGAA
 GAGCTCCGGGACCGGTGAAAGGGTCTCTGCCAAACCTTTCATCGAGACAGTCCCTTCCATAGATGCGC
 TTCCTGTGACATTGGCAATGCAGCTGAATTCTATAAGATTTCCAGCTGGAGATAGGGGAAGTGTATAA
 ACATCCCAATGCCTCTAAAGAGGAAAGGAAGAGATGGCAGGCCAGCTGGACAAACATCTCCGAAAAGG
 ATGAACCTAAAACCAATCATGAGGATGAATGGCAACTTTGCCCGAAGCTTATGACCCAAGAGAGCTGAG
 ACGCAGTTTGTGAGTTAATTCCTTCTGAGGAGAGGCATGAAGCTCTCAGGGAGCTCATGGACCTTACCT
 GAAGATGAAACCCGTGTGGCGCTCTTATGTCCCGCTAAAGAGTGTCCAGAGTCCCTCTGTGAGTACAGT
 TTCAACTCACAGCGTTTCGCGGAACTCCTCCACCAAGTTCAAATATAGATACGAGGGCAAAATCACCA
 ATTACTTTCACAAAACCTTGGCAGATGTCCCTGAAATTATTGAAAGGGATGGCTCTATCGGGGCTGGGC
 AAGTGAGGGAAATGAATCGGGTAAACAGCTGTTTAGACGGTTTCGGAAAAATGAATGCCAGGCAGTCCAAG
 TGCTATGAGATGGAAGATGTCTGAAACATCACTGGCTGTATACTTCAAATACCTCCAGAAGTTTATGA
 ATGCTCATAACCGGTTAAAAGCTCTGGGTTTACCATGAACTCAAAGGAGACCTTAGGGGACCTTTGGG
 CATTGAGGACTCTCTGGAAAGCCAAGATTCAATGGAGTTT

AGCGGACCGACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC
 TGGATTACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:

>MR227208 representing NM_009019
 Red=Cloning site Green=Tags(s)

MAASLPSTLSFSSAPDEIQHPQIKFSEWFKLFRVRSFEKAPEEAQKEKDSSEGKPYLEQSPVVPPEKPGG
 QNSILTQRALKLHPKFSKKFHADGKSSDKAVHQARLRHFCRICGNRFKSDGHSRRYPVHGPVDAKTQSLF
 RKKEKRVTSWPDLIARIFRIDVKADVDSIHPTEFCHDCWSIMHRKFSSSHSQVYFPRKVTVEWHPHTPSC
 DICFTAHRGLKRRHQPNVQLSKLKTVLNHARRDRRRTQARVSSKEVLKKISNCSKIHLSTKLLAVDF
 PAHFVKSISQCICEHILADPVETSCKHLFCRICILRCLKVMGSYCPSCRYPCFPTDLESVPKSFNLILNS
 LMVKCPAQDCNEEVSLEKYNHHVSSHESKETLVHINKGGRPRQHLLSLTRRAQKHRLRELKIQVKEFAD
 KEEGGDVKAVCLTLFLLALARNEHRQADELEAIMQGRGSLQPAVCLAIRVNTFLSCSQYHKMYRTVKA
 ITGRQIFQPLHALRNAEKVLLPGYHPFEWQPPLKNVSSRTDVGIIDGLSGLASSVDEYPVDTIAKRFYD
 SALVLSALMDEEDILEGMRSQDLDDYLNPFVTVVKESCDGMGDVSEKHGSGPAVPEKAVRFSFTVMRIT
 IEHGSQNVKVFEEPKPNSELCKPLCLMLADESDHETLTAILSPLIAEREAMKSELLEMGGIPRTFKF
 IFRGTGYDEKLVREVEGLEASGSVYICTLCDTTRLEASQNLVFHSITRSHAENLQRYEVWRSNPYHESVE
 ELRDRVKVSAKPFIEIVPSIDALHCDIGNAAEFYKIFQLEIGEYVYKHPNASKEERKRWQATLDKHLRKR
 MNLKPIMRMNGNFARKLMTQETVDAVCELIPSEERHEALREMDLYLKMKPVWRSSCPAKECPESLCOYS
 FNSQRFAELLSTKFKYRYEGKITNYFHKTLAHVPEIIERDGSIGAWASEGNESGNKLFRRFRKMNRQSK
 CYEMEDVLKHHWLYTSKYLQKFMNAHNALKSSGF TMNSKETLGDPLGIEDSLESQDSMEF


SGP^{TRTRPLEQKLI SEEDLAANDILDYKDDDDKV}

Chromatograms:

https://cdn.origene.com/chromatograms/mm9039_h10.zip

Restriction Sites:

Sgfl-RsrII

Cloning Scheme:


ACCN: NM_009019

ORF Size: 3120 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
 3. Close the tube and incubate for 10 minutes at room temperature.
 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_009019.2](#), [NP_033045.2](#)

RefSeq Size: 6669 bp

RefSeq ORF: 3123 bp

Locus ID: 19373


UniProt ID: [P15919](#)

Cytogenetics: 2 53.88 cM

MW: 119.6 kDa

Gene Summary: Catalytic component of the RAG complex, a multiprotein complex that mediates the DNA cleavage phase during V(D)J recombination. V(D)J recombination assembles a diverse repertoire of immunoglobulin and T-cell receptor genes in developing B and T-lymphocytes through rearrangement of different V (variable), in some cases D (diversity), and J (joining) gene segments. In the RAG complex, RAG1 mediates the DNA-binding to the conserved recombination signal sequences (RSS) and catalyzes the DNA cleavage activities by introducing a double-strand break between the RSS and the adjacent coding segment. RAG2 is not a catalytic component but is required for all known catalytic activities. DNA cleavage occurs in 2 steps: a first nick is introduced in the top strand immediately upstream of the heptamer, generating a 3'-hydroxyl group that can attack the phosphodiester bond on the opposite strand in a direct transesterification reaction, thereby creating 4 DNA ends: 2 hairpin coding ends and 2 blunt, 5'-phosphorylated ends. The chromatin structure plays an essential role in the V(D)J recombination reactions and the presence of histone H3 trimethylated at 'Lys-4' (H3K4me3) stimulates both the nicking and hairpinning steps. The RAG complex also plays a role in pre-B cell allelic exclusion, a process leading to expression of a single immunoglobulin heavy chain allele to enforce clonality and monospecific recognition by the B-cell antigen receptor (BCR) expressed on individual B-lymphocytes. The introduction of DNA breaks by the RAG complex on one immunoglobulin allele induces ATM-dependent repositioning of the other allele to pericentromeric heterochromatin, preventing accessibility to the RAG complex and recombination of the second allele. In addition to its endonuclease activity, RAG1 also acts as an E3 ubiquitin-protein ligase that mediates monoubiquitination of histone H3. Histone H3 monoubiquitination is required for the joining step of V(D)J recombination. Mediates polyubiquitination of KPNA1.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR227208