

Product datasheet for MR227454L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Trim33 (NM_053170) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Trim33 (NM_053170) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Trim33

Synonyms: 8030451N04Rik; Al413936; Ecto; Tif1g

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_053170

 ORF Size:
 3420 bp

ORF Nucleotide

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(MR227454).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 053170.2, NP 444400.2

 RefSeq Size:
 8868 bp

 RefSeq ORF:
 3423 bp

 Locus ID:
 94093

 UniProt ID:
 Q99PP7

Cytogenetics: 3 F2.2







Gene Summary:

Acts as an E3 ubiquitin-protein ligase. Promotes SMAD4 ubiquitination, nuclear exclusion and degradation via the ubiquitin proteasome pathway (By similarity). May act as a transcriptional repressor (By similarity). Inhibits the transcriptional response to TGF-beta/BMP signaling cascade (By similarity). Plays a role in the control of cell proliferation (By similarity). Its association with SMAD2 and SMAD3 stimulates erythroid differentiation of hematopoietic stem/progenitor. Monoubiquitinates SMAD4 and acts as an inhibitor of SMAD4-dependent TGF-beta/BMP signaling cascade (Monoubiquitination of SMAD4 hampers its ability to form a stable complex with activated SMAD2/3 resulting in inhibition of TGF-beta/BMP signaling cascade) (By similarity).[UniProtKB/Swiss-Prot Function]