

Product datasheet for MR227499L4

Havcr2 (NM_134250) Mouse Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Havcr2 (NM_134250) Mouse Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Havcr2
Synonyms:	TIM-3; Tim3; Timd3
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR227499).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.

ACCN:	NM_134250
ORF Size:	843 bp



[View online »](#)

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_134250.2](#)

RefSeq Size: 2725 bp

RefSeq ORF: 846 bp

Locus ID: 171285


UniProt ID: [Q8VIM0](#)

Cytogenetics: 11 B1.1


Gene Summary:

Cell surface receptor implicated in modulating innate and adaptive immune responses. Generally accepted to have an inhibiting function. Reports on stimulating functions suggest that the activity may be influenced by the cellular context and/or the respective ligand (PubMed:18006747). Regulates macrophage activation (PubMed:11823861). Inhibits T-helper type 1 lymphocyte (Th1)-mediated auto- and alloimmune responses and promotes immunological tolerance (PubMed:14556006, PubMed:18006747). In CD8+ cells attenuates TCR-induced signaling, specifically by blocking NF-kappaB and NFAT promoter activities resulting in the loss of IL-2 secretion. The function may implicate its association with LCK proposed to impair phosphorylation of TCR subunits (By similarity). In contrast, shown to activate TCR-induced signaling in T-cells probably implicating ZAP70, LCP2, LCK and FYN (PubMed:21807895). Expressed on Treg cells can inhibit Th17 cell responses (By similarity). Receptor for LGALS9. Binding to LGALS9 is believed to result in suppression of T-cell responses; the resulting apoptosis of antigen-specific cells may implicate HAVCR2 phosphorylation and disruption of its association with BAG6 (PubMed:22863785). Binding to LGALS9 is proposed to be involved in innate immune response to intracellular pathogens. Expressed on Th1 cells interacts with LGALS9 expressed on Mycobacterium tuberculosis-infected macrophages to stimulate antibactericidal activity including IL-1 beta secretion and to restrict intracellular bacterial growth (PubMed:20937702). However, the function as receptor for LGALS9 has been challenged (By similarity). Also reported to enhance CD8+ T-cell responses to an acute infection such as by Listeria monocytogenes (PubMed:24567532). Receptor for phosphatidylserine (PtSer); PtSer-binding is calcium-dependent (PubMed:20083673). May recognize PtSer on apoptotic cells leading to their phagocytosis. Mediates the engulfment of apoptotic cells by dendritic cells (PubMed:19224762). Expressed on T-cells, promotes conjugation but not engulfment of apoptotic cells (PubMed:20083673). Expressed on dendritic cells (DCs) positively regulates innate immune response and in synergy with Toll-like receptors promotes secretion of TNF-alpha (PubMed:18006747). In tumor-infiltrating DCs suppresses nucleic acid-mediated innate immune response by interaction with HMGB1 and interfering with nucleic acid-sensing and trafficking of nucleic acids to endosomes (PubMed:22842346). Can enhance mast cell production of Th2 cytokines IL-4, IL-6 and IL-13 (PubMed:17620455). Expressed on natural killer (NK) cells acts as a coreceptor to enhance IFN-gamma production in response to LGALS9. In contrast, shown to suppress NK cell-mediated cytotoxicity (By similarity). Negatively regulates NK cell function in LPS-induced endotoxic shock (PubMed:25337993).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR227499L4



Double digestion of MR227499L4 using SgfI and MluI