

Product datasheet for MR227518

Sirt6 (NM_001163430) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Sirt6 (NM_001163430) Mouse Tagged ORF Clone
Tag: Myc-DDK
Symbol: Sirt6
Synonyms: 2810449N18Rik; AI043036; Sir2l6
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
ORF Nucleotide Sequence: >MR227518 representing NM_001163430
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCGCGATCGCC

ATGTGGCAGTCTCCAGCGTGGTTTTCCACACGGGCGCCGGCATCAGCACCGCTCTGGCATCCCCGACT
 TCAGAGGCCCCCATGGCGTGTGGACCATGGAGGAACGCGGCTGGCCCCAAGTTTGACACCACCTTCGA
 GAATGCTCGGCCCTCGAAGACCCACATGGCCCTGGTTCAGCTAGAACGCATGGGCTTCTCAGCTTCTG
 GTCAGCCAGAACGTAGACGGGCTGCACGTGCGCTCGGGCTTCCCCAGGGACAAGCTGGCAGAGCTGCACG
 GAAACATGTTGTAGAGGAATGTCCCAAGTGAAGACGCAGTACGTGAGACACGGTTGTGGGCACCAT
 GGGCTCAAGGCCACAGGCCGGCTCTGCACCGTGGCCAAGACCAGGGGACTTCGGGCCTGTAGAGGGGAG
 CTGAGAGACACCATCTGGACTGGGAGGACTCGTTGCCTGACCGGGACCTGATGCTCGCTGATGAGGCCA
 GCAGGACCGCAGACCTGTCTGTCAACCCTGGGTACCTCGCTGCAGATCCGCCCCAGTGGGAACCTGCCCT
 TGCCACTAAGCGCCGAGGAGGCCGTCTGGTCATTGTCAACCTGCAACCCACAAAACATGACCGCCAGGT
 GACCTGCGCATCCACGGTACGTGGATGAGGTGATGTGCAGACTCATGAAGCATCTGGGGCTGGAGATTC
 CAGCCTGGGATGGACCTGCGTGTAGACAAAGCCCTGCCACCTCTGCCTCGCCAGTAGCACTCAAGGC
 TGAGCCCCCGTGCATCTCAATGGTGCAGTGCATGTTTCGTATAAGTCCAAGCCCAACAGCCCTATACTC
 CACAGGCCCCCCAAAAGAGTGAAGACCGAGGCTGCCCCCAGC

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

Protein Sequence: >MR227518 representing NM_001163430
 Red=Cloning site Green=Tags(s)


MWQSSSVFHTGAGISTASGIPDFRGPVHVTMEERGLAPKFDTTFENARPSKTHMALVQLERMGFLSFL
 VSQNVDDLHVRSGFPRDKLAELHGNMFVEECPKCKTQYVRDVTVGTMLKATGRLCTVAKTRGLRACRGE
 LRDTILDWEDSLPDRDLMLADEASRTADLSVTLGTSLQIRPSGNLPLATKRRGGRLVIVNLQPTKHDRQA
 DLRIHGYVDEVMCRLMKHLGLEIPAWDGPCVLDKALPPLPRPVALKAEPVHVLNGAVHVSYSKSKPNPIL
 HRPPKRVKTEAAPS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV


Restriction Sites:

Sgfl-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_001163430

ORF Size: 882 bp

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001163430.2
RefSeq Size:	1668 bp
RefSeq ORF:	885 bp
Locus ID:	50721
Cytogenetics:	10 39.72 cM
MW:	32.9 kDa
Gene Summary:	NAD-dependent protein deacetylase. Has deacetylase activity towards histone H3K9Ac and H3K56Ac. Modulates acetylation of histone H3 in telomeric chromatin during the S-phase of the cell cycle. Deacetylates histone H3K9Ac at NF-kappa-B target promoters and may down-regulate the expression of a subset of NF-kappa-B target genes. Deacetylation of nucleosomes interferes with RELA binding to target DNA. May be required for the association of WRN with telomeres during S-phase and for normal telomere maintenance. On DNA damage, promotes DNA end resection via deacetylation of RBBP8. Has very weak deacetylase activity and can bind NAD(+) in the absence of acetylated substrate (By similarity). Acts as a corepressor of the transcription factor Hif1a to control the expression of multiple glycolytic genes to regulate glucose homeostasis. Required for genomic stability. Required for normal IGF1 serum levels and normal glucose homeostasis. Modulates cellular senescence and apoptosis. Regulates the production of TNF protein. Has a role in the regulation of life span in male mice, but not in female mice.[UniProtKB/Swiss-Prot Function]