

Product datasheet for MR227522L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Nr1h3 (NM_013839) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Nr1h3 (NM 013839) Mouse Tagged ORF Clone Lentiviral Particle

Symbol: Nr1h3

Synonyms: AU018371; LXR; RLD1; Unr1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_013839 **ORF Size:** 1335 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(MR227522).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 013839.4, NP 038867.2

 RefSeq Size:
 1955 bp

 RefSeq ORF:
 1338 bp

 Locus ID:
 22259

 UniProt ID:
 Q9Z0Y9

 Cytogenetics:
 2 50.52 cM







Gene Summary:

Nuclear receptor that exhibits a ligand-dependent transcriptional activation activity (PubMed:18055760, PubMed:19520913, PubMed:20427281). Interaction with retinoic acid receptor (RXR) shifts RXR from its role as a silent DNA-binding partner to an active ligand-binding subunit in mediating retinoid responses through target genes defined by LXRES. LXRES are DR4-type response elements characterized by direct repeats of two similar hexanuclotide half-sites spaced by four nucleotides. Plays an important role in the regulation of cholesterol homeostasis, regulating cholesterol uptake through MYLIP-dependent ubiquitination of LDLR, VLDLR and LRP8. Interplays functionally with RORA for the regulation of genes involved in liver metabolism.[UniProtKB/Swiss-Prot Function]