

Product datasheet for **MR227544L1V**

Tmem173 (NM_028261) Mouse Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Tmem173 (NM_028261) Mouse Tagged ORF Clone Lentiviral Particle
Symbol:	Tmem173
Synonyms:	2610307O08Rik; ERIS; Mita; MPYS; STING
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_028261
ORF Size:	1134 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR227544).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_028261.1 , NP_082537.1
RefSeq Size:	2302 bp
RefSeq ORF:	1137 bp
Locus ID:	72512
UniProt ID:	Q3TBT3
Cytogenetics:	18



[View online »](#)

Gene Summary:

Facilitator of innate immune signaling that acts as a sensor of cytosolic DNA from bacteria and viruses and promotes the production of type I interferon (IFN-alpha and IFN-beta) (PubMed:18818105, PubMed:19433799, PubMed:19776740, PubMed:26229117, PubMed:26669264). Innate immune response is triggered in response to non-CpG double-stranded DNA from viruses and bacteria delivered to the cytoplasm (PubMed:18818105, PubMed:19433799, PubMed:19776740, PubMed:26229117, PubMed:26669264). Acts by binding cyclic dinucleotides: recognizes and binds cyclic di-GMP (c-di-GMP), a second messenger produced by bacteria, and cyclic GMP-AMP (cGAMP), a messenger produced by CGAS in response to DNA virus in the cytosol (PubMed:21947006, PubMed:23722158, PubMed:23258412, PubMed:23519410, PubMed:23910378). Upon binding of c-di-GMP or cGAMP, TMEM173/STING oligomerizes, translocates from the endoplasmic reticulum and is phosphorylated by TBK1 on the pLxIS motif, leading to recruitment and subsequent activation of the transcription factor IRF3 to induce expression of type I interferon and exert a potent anti-viral state (PubMed:25636800). In addition to promote the production of type I interferons, plays a direct role in autophagy (PubMed:30568238). Following cGAMP-binding, TMEM173/STING buds from the endoplasmic reticulum into COPII vesicles, which then form the endoplasmic reticulum-Golgi intermediate compartment (ERGIC) (By similarity). The ERGIC serves as the membrane source for WIPI2 recruitment and LC3 lipidation, leading to formation of autophagosomes that target cytosolic DNA or DNA viruses for degradation by the lysosome (By similarity). The autophagy- and interferon-inducing activities can be uncoupled and autophagy induction is independent of TBK1 phosphorylation (By similarity). Autophagy is also triggered upon infection by bacteria: following c-di-GMP-binding, which is produced by live Gram-positive bacteria, promotes reticulophagy (PubMed:29056340). Exhibits 2',3' phosphodiester linkage-specific ligand recognition: can bind both 2'-3' linked cGAMP (2'-3'-cGAMP) and 3'-3' linked cGAMP but is preferentially activated by 2'-3' linked cGAMP (PubMed:26300263). The preference for 2'-3'-cGAMP, compared to other linkage isomers is probably due to the ligand itself, which adopts an organized free-ligand conformation that resembles the TMEM173/STING-bound conformation and pays low energy costs in changing into the active conformation (By similarity). May be involved in translocon function, the translocon possibly being able to influence the induction of type I interferons (By similarity). May be involved in transduction of apoptotic signals via its association with the major histocompatibility complex class II (MHC-II) (PubMed:18559423).[UniProtKB/Swiss-Prot Function]