

## Product datasheet for MR227555L4

## Xiap (NM\_009688) Mouse Tagged Lenti ORF Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Xiap (NM\_009688) Mouse Tagged Lenti ORF Clone

Tag: mGFP Symbol: Xiap

Synonyms: 1110015C02Rik; A; Aipa; Api3; Bir; Birc4; I; IAP3; IL; ILP-1; MIHA

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)


E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(MR227555).

Sequence:

Restriction Sites: Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_009688

ORF Size: 1488 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com **OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 009688.2</u>, <u>NP 033818.2</u>

RefSeq Size: 6550 bp
RefSeq ORF: 1491 bp
Locus ID: 11798
UniProt ID: Q60989
Cytogenetics: X A4


**Gene Summary:** The protein encoded by this gene is a member of the inhibitor of apoptosis (IAP) family of

proteins. While first identified for its role in blocking apoptosis, this protein modulates many other signaling processes including nuclear factor kappa-light-chain-enhancer of activated B cells (NF-kB) pathways and inflammatory responses. This protein blocks apoptosis by binding and inhibiting target caspases after they have been activated. Binding occurs to some, but not all, caspases. This protein has several conserved regions, including baculoviral IAP repeat (BIR) motifs and a RING finger E3 ligase domain. In humans, mutations in this gene are linked to immunodeficiency in X-linked lymphoproliferative syndrome type-2 (XLP-2). A pseudogene of this gene is found on chromosome 7. Alternative splicing results in multiple transcript

variants. [provided by RefSeq, Aug 2014]



## **Product images:**



Circular map for MR227555L4