

Product datasheet for MR227573L3


Prx (NM_019412) Mouse Tagged Lenti ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	Prx (NM_019412) Mouse Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Prx
Synonyms:	L-Periaxin
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(MR227573).
Restriction Sites:	SgfI-RsrII
Cloning Scheme:	
	Cloning sites used for ORF Shuttling:
	$\frac{SgfI}{GCGATCGC} = \frac{ORF}{ATG} = \frac{RsrII}{CGGACCG} = \frac{SgfI}{CGGACCG} = \frac{SgfI}{CGGACC$



* The last codon before the Stop codon of the ORF.

ACCN: ORF Size: NM_019412

444 bp



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

ORIGENE Prx (NM_019412) Mouse Tagged Lenti ORF Clone – MR227573L3


OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.
	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 019412.2, NP 062285.1</u>
RefSeq Size:	5260 bp
RefSeq ORF:	447 bp
Locus ID:	19153
UniProt ID:	<u>055103</u>
Cytogenetics:	7 15.91 cM

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Section 2019 Prx (NM_019412) Mouse Tagged Lenti ORF Clone – MR227573L3

Scaffolding protein that functions as part of a dystroglycan complex in Schwann cells, and as Gene Summary: part of EZR and AHNAK-containing complexes in eye lens fiber cells (PubMed:11430802, PubMed:21745462, PubMed:22764250). Required for the maintenance of the peripheral myelin sheath that is essential for normal transmission of nerve impulses and normal perception of sensory stimuli (PubMed:10839370). Required for normal transport of MBP mRNA from the perinuclear to the paranodal regions (PubMed:15356632). Required for normal remyelination after nerve injury (PubMed:10839370). Required for normal elongation of Schwann cells and normal length of the internodes between the nodes of Ranvier. The demyelinated nodes of Ranvier permit saltatory transmission of nerve impulses; shorter internodes cause slower transmission of nerve impulses (PubMed:15356632, PubMed:23022068). Required for the formation of appositions between the abaxonal surface of the myelin sheath and the Schwann cell plasma membrane; the Schwann cell cytoplasm is restricted to regions between these appositions (PubMed:15356632, PubMed:23022068). Required for the formation of Cajal bands and of Schmidt-Lanterman incisures that correspond to short, cytoplasm-filled regions on myelinated nerves (PubMed:23022068, PubMed:22764250). Recruits DRP2 to the Schwann cell plasma membrane (PubMed:11430802, PubMed:23022068, PubMed:22764250). Required for normal protein composition of the eye lens fiber cell plasma membrane and normal eye lens fiber cell morphology (PubMed:21745462).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR227573L3

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US