

Product datasheet for **MR227663**

Cd36 (NM_007643) Mouse Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Cd36 (NM_007643) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Cd36
Synonyms:	FAT; GPIV; Scarb3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

ORF Nucleotide Sequence:

>MR227663 ORF sequence
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGCATCGCC**

ATGGGCTGTGATCGGAAGTGTGGGCTCATTGCTGGAGCTGTTATTGGTGCAGTCCTGGCTGTGTTTGGAG
 GCATTCTCATGCCAGTCGGAGACATGCTTATTGAGAAGACAATCAAAGGGAAGTTGTCCTTGAAGAAGG
 AACCACTGCTTTCAAAAAGTGGGTTAAAACAGGCACCACTGTGTACAGACAGTTTTGGATCTTTGATGTG
 CAAAACCCAGATGACGTGGCAAAGAACAGCAGCAAATCAAGGTTAAACAAAGAGGTCCTTACACATACA
 GAGTTCGTTATCTAGCCAAGGAAAATAAAGTCAAGGACCCCGAGGACCACACTGTGTCTTTGTACAGCC
 CAATGGAGCCATCTTTGAGCCTTCACTGTCTGTTGGAACAGAGGATGACAACCTTACAGTCTGAATCTG
 GCTGTAGCAGCTGCACCACATCTACCAAAATTCATTTGTTCAAGTTGTGCTCAACTCTTTATAAAAA
 AGTCCAAGTCTTCTATGTTCCAAACAAGATCTTTGAAAGAAGTCTGTGGGGTTACAAAGATCCATTCT
 CAGTTTGGTTCCATATCCTATAAGTACCACAGTTGGTGTGTTTTATCCTTACAATGACACTGTAGATGGA
 GTTTATAAAGTTTTCAATGGAAAGGATAACATAAGCAAAGTTGCCATAATTGAGTCTATAAAGGAAAA
 GGAATTTGTCCTATTGGCCAAGCTATTGCGACATGATTAAATGGCAGACGACGAGCCTCCTTTCCACCTTT
 TGTGAAAAGTCTCGGACATTGAGATCTTTTCTCTGACATTTGCGAGTCTATCTACGCTGTGTTCCGGA
 TCTGAAATCGACCTAAAGGAATCCCGGTGACAGATTTGTTCTTCCAGCCAATGCCTTTGCATCACCCC
 TCCAGAATCCAGACAACCATTTCTGCACTGAAAAAGTAACTCCAATAACTGTACATCTTATGGTGT
 GCTAGACATTTGCAAATGCAAAGAAGGAAAGCCTGTGTATTTTCGCTCCACATTTCTACATGCAAGT
 CCAGATGTTTCCAGAACCTATTGAAGGCTTACATCCAATGAAGATGAGCATAGGACATACTTAGATGTGG
 AACCCATAACTGGATTCACCTACAATTTGCAAAACGACTGCAGTCAACATATTGGTCAAGCCAGCTAG
 AAAAAATAGAAGCATTAAAGAATCTGAAGAGACCTTACATTGTACCTATACTGTGGCTAAAATGAGACTGGG
 ACCATTGGTGATGAAAAAGCAGAAATGTTCAAAACACAAGTACTGGGAAAATCAAGCTCCTTGGCATGG
 TAGAGATGGCCTTACTTGGGATTGGAGTGGTGTGTTTGTGCTTTTATGATTTCAATTGTGCTTGCAA
 ATCCAAGAATGGAAAA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence:

>MR227663 protein sequence
 Red=Cloning site Green=Tags(s)

MGCDRNCGLIAGAVIGAVLAVFGGILMPVGDMLIEKTIKREVVLEEGTTAFKNWVKTGTTVYRQFWIFDV
 QNPDDVAKNSSKIKVKQRPYTYRVRYLAKENITQDPEDHTVSFVQPNGAIFEPSLSVGTEDDNFTVLNL
 AVAAAPHIYQNSFVQVVLNSLIKKSMMFQTRSLKELLWGYKDPFLSLVPYPISTTVGVFYPYNDTVDG
 VYKVFNGKDNISKVAIIIESYKGRNLSYWPSYCDMINGTDAASFPPFVEKSRTRLRFFSSDICRSIYAVFG
 SEIDLKGI PVYRFVLPANAFASPLQNPDNHCFCTEKVISNNCTS YGVLDIGKCKEKPVIYISLPHFLHAS
 PDVSEPIEGLHPNEDEHRTYLDVEPITGFTLQFAKRLQVNILVKPARKIEALKNLKRPYIVPILWLNETH
 TIGDEKAEMFKTQVTGKIKLLGMVEMALLGIGVVMFVAFMISYACKSKNGK

TRTRPLEQ**KL**ISEEDLAANDILDYKDDDDKV

Restriction Sites:

SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shutting:



* The last codon before the Stop codon of the ORF

ACCN: NM_007643

ORF Size: 1416 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_007643.4](#), [NP_031669.3](#)
RefSeq Size: 3542 bp

RefSeq ORF: 1419 bp

Locus ID: 12491


UniProt ID: [Q08857](#)

Cytogenetics: 5 8.11 cM

MW: 52.7 kDa

Gene Summary: Multifunctional glycoprotein that acts as receptor for a broad range of ligands. Ligands can be of proteinaceous nature like thrombospondin, fibronectin, collagen or amyloid-beta as well as of lipidic nature such as oxidized low-density lipoprotein (oxLDL), anionic phospholipids, long-chain fatty acids and bacterial diacylated lipopeptides (PubMed:7685021). They are generally multivalent and can therefore engage multiple receptors simultaneously, the resulting formation of CD36 clusters initiates signal transduction and internalization of receptor-ligand complexes. The dependency on coreceptor signaling is strongly ligand specific. Cellular responses to these ligands are involved in angiogenesis, inflammatory response, fatty acid metabolism, taste and dietary fat processing in the intestine (Probable) (PubMed:19847289, PubMed:20037584, PubMed:23395392). Binds long-chain fatty acids and facilitates their transport into cells, thus participating in muscle lipid utilization, adipose energy storage, and gut fat absorption (By similarity). In the small intestine, plays a role in proximal absorption of dietary fatty acid and cholesterol for optimal chylomicron formation, possibly through the activation of MAPK1/3 (ERK1/2) signaling pathway (By similarity) (PubMed:17507371, PubMed:18753675, PubMed:21610069). Involved in oral fat perception and preferences (PubMed:16276419). Detection into the tongue of long-chain fatty acids leads to a rapid and sustained rise in flux and protein content of pancreatobiliary secretions (By similarity) (PubMed:16276419). In taste receptor cells, mediates the induction of an increase in intracellular calcium levels by long-chain fatty acids, leading to the activation of the gustatory neurons in the nucleus of the solitary tract (PubMed:18162488). Important factor in both ventromedial hypothalamus neuronal sensing of long-chain fatty acid and the regulation of energy and glucose homeostasis (By similarity) (PubMed:23557700). Receptor for thrombospondins, THBS1 and THBS2, mediating their antiangiogenic effects (PubMed:15748999). As a coreceptor for TLR4:TLR6 heterodimer, promotes inflammation in monocytes/macrophages. Upon ligand binding, such as oxLDL or amyloid-beta 42, interacts with the heterodimer TLR4:TLR6, the complex is internalized and triggers inflammatory response, leading to NF-kappa-B-dependent production of CXCL1, CXCL2 and CCL9 cytokines, via MYD88 signaling pathway, and CCL5 cytokine, via TICAM1 signaling pathway, as well as IL1B secretion, through the priming and activation of the NLRP3 inflammasome (PubMed:20037584, PubMed:23812099). Selective and nonredundant sensor of microbial diacylated lipopeptide that signal via TLR2:TLR6 heterodimer, this cluster triggers signaling from the cell surface, leading to the NF-kappa-B-dependent production of TNF, via MYD88 signaling pathway and subsequently is targeted to the Golgi in a lipid-raft dependent pathway (By similarity) (PubMed:15690042, PubMed:19847289).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR227663