

Product datasheet for MR227669

Cd3e (NM 007648) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Cd3e (NM_007648) Mouse Tagged ORF Clone

Tag: Myc-DDK

Symbol: Cd3e

Synonyms: Al504783; CD3; CD3epsilon; T3e

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >MR227669 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGCGGTGGAACACTTTCTGGGGCATCCTGTGCCTCAGCCTCCTAGCTGTTGGCACTTGCCAGGACGATG CCGAGAACATTGAATACAAAGTCTCCATCTCAGGAACCAGTGTAGAGTTGACGTGCCCTCTAGACAGTGA CGAGAACTTAAAATGGGAAAAAAATGGCCAAGAGCTGCCTCAGAAGCATGATAAGCACCTGGTGCTCCAG GATTTCTCGGAAGTCGAGGACAGTGGCTACTACGTCTGCTACACCACCAGCCTCAAATAAAAACACGTACT TGTACCTGAAAGCTCGAGTGTGTGAGTACTGTGTGGAGGTGGACCTGACAGCAGTAGCCATAATCATCAT TGTTGACATCTGTATCACTCTGGGCTTGCTGATGGTCATTTATTACTGGAGCAAGAATAGGAAGGCCAAG GCCAAGCCTGTGACCCGAGGAACCGGTGCTGGTAGCAGGCCCAGAGGGCAAAACAAGGAGCGGCCACCAC CTGTTCCCAACCCAGACTATGAGCCCATCCGCAAAGGCCAGCGGGACCTGTATTCTGGCCTGAATCAGAG

AGCAGTC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR227669 protein sequence

Red=Cloning site Green=Tags(s)

MRWNTFWGILCLSLLAVGTCQDDAENIEYKVSISGTSVELTCPLDSDENLKWEKNGQELPQKHDKHLVLQ DFSEVEDSGYYVCYTPASNKNTYLYLKARVCEYCVEVDLTAVAIIIIVDICITLGLLMVIYYWSKNRKAK

AKPVTRGTGAGSRPRGQNKERPPPVPNPDYEPIRKGQRDLYSGLNQRAV

TRTRPLEQKLISEEDLAANDILDYKDDDDKV



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com ORÏGENE

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_007648

ORF Size: 567 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 007648.5</u>

RefSeq Size: 1436 bp RefSeq ORF: 570 bp



Locus ID: 12501 UniProt ID: <u>P22646</u>

 Cytogenetics:
 9 24.84 cM

 MW:
 21.4 kDa

Gene Summary: Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role


in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways. In addition of this role of signal transduction in

T-cell activation, CD3E plays an essential role in correct T-cell development

(PubMed:19956738, PubMed:24899501). Participates also in internalization and cell surface down-regulation of TCR-CD3 complexes via endocytosis sequences present in CD3E cytosolic

region.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR227669