

# Product datasheet for MR227915

# Lair1 (NM\_001302677) Mouse Tagged ORF Clone

# **Product data:**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	Lair1 (NM_001302677) Mouse Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Lair1
Synonyms:	5133400O11Rik; BB115266; D7Bwg0421e; Lair-1
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin
ORF Nucleotide Sequence:	>MR227915 representing NM_001302677 Red=Cloning site Blue=ORF Green=Tags(s)
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGTCACTTCATCCAGTTATCCTGCTGGTGCTTGTGCTGTGCCTGGGATGGAAAATTAACACACAGGAGG GACTCCCAAACAACAAAAGACAGCAGCAGAGAGGCCAGAAGAGAGGGCTAAATCTAGCTACTAATGGCCTGGA GATGACTCCAGACATAGTTGCAGATGACAGGCTTCCTGAGGACAGATGGACAGAAACCTGGACCCCAGTT GCAGGAGACCTTCAAGAGGTGACGTATATCCAGCTGGACCATCACTCCCTCACACAGAGGGCAGTCGGAG CTGTGACCTCACAGAGCACAGATATGGCTGAGTCCAGCACATATGCAGCCATCATCAGACAC
	ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAG <b>GTTTAA</b>
Protein Sequence:	>MR227915 representing NM_001302677 <mark>Red</mark> =Cloning site Green=Tags(s)
	MSLHPVILLVLVLCLGWKINTQEGLPNNKRQQQRPEERLNLATNGLEMTPDIVADDRLPEDRWTETWTPV AGDLQEVTYIQLDHHSLTQRAVGAVTSQSTDMAESSTYAAIIRH
	TRTRPLEQKLISEEDLAANDILDYKDDDDKV
Restriction Sites:	Sgfl-Mlul



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US




#### **Cloning Scheme:**



\* The last codon before the Stop codon of the ORF

## Plasmid Map:



ACCN:	
ORF Size:	
OTI Disclaimer:	

## NM\_001302677

## 342 bp

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

<b>ORIGENE</b> Lair1 (I	NM_001302677) Mouse Tagged ORF Clone – MR227915
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
RefSeq:	<u>NM 001302677.1, NP 001289606.1</u>
RefSeq Size:	3065 bp
RefSeq ORF:	345 bp
Locus ID:	52855
UniProt ID:	<u>Q8BG84</u>
Cytogenetics:	7 2.31 cM
MW:	13.3 kDa
Gene Summary:	Functions as an inhibitory receptor that plays a constitutive negative regulatory role on cytolytic function of natural killer (NK) cells, B-cells and T-cells. Activation by Tyr phosphorylation results in recruitment and activation of the phosphatases PTPN6 and PTPN11. It also reduces the increase of intracellular calcium evoked by B-cell receptor ligation. May also play its inhibitory role independently of SH2-containing phosphatases. Modulates cytokine production in CD4+ T-cells, down-regulating IL2 and IFNG production while inducing secretion of transforming growth factor beta. Down-regulates also IgG and IgE production in B-cells as well as IL8, IL10 and TNF secretion. Inhibits proliferation and induces apoptosis in myeloid leukemia cell lines as well as prevents nuclear translocation of NF-kappa-B p65 subunit/RELA and phosphorylation of I-kappa-B alpha/CHUK in these cells. Inhibits the differentiation of peripheral blood precursors towards dendritic cells (By similarity).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US