

## Product datasheet for MR228024

### FasI (NM\_001205243) Mouse Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** FasI (NM\_001205243) Mouse Tagged ORF Clone  
**Tag:** Myc-DDK  
**Symbol:** FasI  
**Synonyms:** APT1LG1; CD95-L; CD95L; CD178; Fas-L; Faslg; gld; Tnfsf6; Tnlg1a  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Restriction Sites:** SgfI-MluI  
**Cloning Scheme:**

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF


**ACCN:** NM\_001205243  
**ORF Size:** 393 bp



[View online >](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<u>NM_001205243.1, NP_001192172.1</u>
<b>RefSeq Size:</b>	1889 bp
<b>RefSeq ORF:</b>	396 bp
<b>Locus ID:</b>	14103
<b>Cytogenetics:</b>	1 69.95 cM
<b>MW:</b>	15 kDa
<b>Gene Summary:</b>	Cytokine that binds to TNFRSF6/FAS, a receptor that transduces the apoptotic signal into cells (PubMed:7511063). Involved in cytotoxic T-cell-mediated apoptosis, natural killer cell-mediated apoptosis and in T-cell development (PubMed:19794494, PubMed:7532682). Initiates fratricidal/suicidal activation-induced cell death (AICD) in antigen-activated T-cells contributing to the termination of immune responses (PubMed:19794494). TNFRSF6/FAS-mediated apoptosis has also a role in the induction of peripheral tolerance (PubMed:10779162). Binds to TNFRSF6B/DcR3, a decoy receptor that blocks apoptosis (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for MR228024