

Product datasheet for **MR228081**

Cd84 (NM_001252472) Mouse Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Cd84 (NM_001252472) Mouse Tagged ORF Clone
Tag: Myc-DDK
Symbol: Cd84
Synonyms: A130013D22Rik; CDw84; SLAMF5
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
ORF Nucleotide Sequence: >MR228081 representing NM_001252472
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGCCCCAGCGCCATCTGTGGATCTGGTTCCTTTGCCTACAAACCTGGTCTGAAGCAGCAGGAAAAGATG
CAGACCCGGTGGTAATGAATGGGATTCTGGGGAGTCAGTTACTTTCCTCTTAAATATTCAAGAACCAA
GAAAATTGACAACATTGCCTGGACTTCTCAATCATCTGTTGCTTTATAAAACCAGGAGTCAATAAAGCT
GAAGTTACCATAACCCAGGGCACTTATAAAGGACGAATAGAAATCATAGATCAGAAGTATGACCTGGTCA
TTAGAGACCTGAGGATGGAAGATGCAGGAATTACAAAGCAGACATCAATGAAGAGAATGAGGAAACCAT
CACCAAGATCTACTACCTTCATATCTACCGTAAGTTATGGCAGCACGGGGCCTTGGATTTACTTTTGATT

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR228081 representing NM_001252472
Red=Cloning site Green=Tags(s)

MAQRHLWIWFLCLQTWSEAAGKDADPVVMNGILGESVTFLLNIQEPKKIDNIAWTSQSSVAFIKPGVNKA
EVTITQGTYKGRIEIIDQKYDLVIRDLRMEAGTYKADINEENEETITKIYYLHIYRKLWQHGDLLLI

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: Sgfl-MluI




[View online »](#)

Cloning Scheme:



Plasmid Map:



ACCN: NM_001252472

ORF Size: 420 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001252472.1</u> , <u>NP_001239401.1</u>
RefSeq Size:	3520 bp
RefSeq ORF:	423 bp
Locus ID:	12523
UniProt ID:	<u>Q18PI6</u>
Cytogenetics:	1 79.54 cM
MW:	16.6 kDa
Gene Summary:	Self-ligand receptor of the signaling lymphocytic activation molecule (SLAM) family. SLAM receptors triggered by homo- or heterotypic cell-cell interactions are modulating the activation and differentiation of a wide variety of immune cells and thus are involved in the regulation and interconnection of both innate and adaptive immune response. Activities are controlled by presence or absence of small cytoplasmic adapter proteins, SH2D1A/SAP and/or SH2D1B/EAT-2 (PubMed:20962259). Can mediate natural killer (NK) cell cytotoxicity dependent on SH2D1A and SH2D1B (PubMed:20962259). Increases proliferative responses of activated T-cells and SH2D1A/SAP does not seem to be required for this process. Homophilic interactions enhance interferon gamma/IFNG secretion in lymphocytes and induce platelet stimulation via a SH2D1A/SAP-dependent pathway. May serve as a marker for hematopoietic progenitor cells (By similarity). Required for a prolonged T-cell:B-cell contact, optimal T follicular helper function, and germinal center formation (PubMed:20153220). In germinal centers involved in maintaining B cell tolerance and in preventing autoimmunity (PubMed:25801429). In mast cells negatively regulates high affinity immunoglobulin epsilon receptor signaling; independent of SH2D1A and SH2D1B but implicating FES and PTPN6/SHP-1 (By similarity). In macrophages enhances LPS-induced MAPK phosphorylation and NF-kappaB activation and modulates LPS-induced cytokine secretion; involving ITSM 2 (PubMed:20628063). Positively regulates macroautophagy in primary dendritic cells via stabilization of IRF8; inhibits TRIM21-mediated proteasomal degradation of IRF8 (By similarity).[UniProtKB/Swiss-Prot Function]