

## **Product datasheet for MR228307**

## Psma7 (NM 001289476) Mouse Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Psma7 (NM\_001289476) Mouse Tagged ORF Clone

Tag: Myc-DDK
Symbol: Psma7

Synonyms: C6-I

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

ORF Nucleotide >MR228307 representing NM\_001289476
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGCCTTTGCAGGTCTCACCGCTGATGCAAGGATAGTCATCAACAGAGCCCGGGTAGAGTGCCAGAGCC
ACCGGCTGACAGTGGAGGACCCAGTGACTGTGGAGTACATCACCCGCTACATTGCGAGTCTGAAGCAGCG
TTATACACAGAGCAATGGGCGCAGGCCATTTGGTATCTCGGCCCTAATTGTGGGTTTTGACTTTGATGGC
ACTCCCAGACTCTATCAGACTGACCCCTCGGGCACATACCATGCTTGGAAGGCCAATGCCATAGGCCGGG
GCGCCAAGTCAGTGCGTGAATTTCTGGAGAAGAACATACACAGATGATGCCATTGAAACAGATGATCTGAC
CATCAAACTTGTGATCAAGGCACTGTTAGAGGTGGTCCAGTCAGGTGGCAAAAACATCGAACTTGCCGTC
ATGAGGCGGGATCAGCCGCTCAAGATTCTAAATCCTGAAGAAATTGAGAAGTATGTTGCTGAAATTGAGA
AGGAGAAAGAAAATGAAAAGAAGAAGCAAAAGAAAGCATCT

 ${\color{blue} \textbf{ACGCGT}} \textbf{ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT}$ 

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR228307 representing NM\_001289476

Red=Cloning site Green=Tags(s)

MAFAGLTADARIVINRARVECQSHRLTVEDPVTVEYITRYIASLKQRYTQSNGRRPFGISALIVGFDFDG TPRLYQTDPSGTYHAWKANAIGRGAKSVREFLEKNYTDDAIETDDLTIKLVIKALLEVVQSGGKNIELAV

MRRDQPLKILNPEEIEKYVAEIEKEKEENEKKKQKKAS

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:** Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




## **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

## Plasmid Map:



ACCN: NM\_001289476

ORF Size: 534 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.



MW:

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 001289476.1, NP 001276405.1

20.6 kDa

RefSeq Size:1591 bpRefSeq ORF:537 bpLocus ID:26444Cytogenetics:2 H4

**Gene Summary:** Component of the 20S core proteasome complex involved in the proteolytic degradation of

most intracellular proteins. This complex plays numerous essential roles within the cell by associating with different regulatory particles. Associated with two 19S regulatory particles, forms the 26S proteasome and thus participates in the ATP-dependent degradation of ubiquitinated proteins. The 26S proteasome plays a key role in the maintenance of protein

homeostasis by removing misfolded or damaged proteins that could impair cellular

functions, and by removing proteins whose functions are no longer required. Associated with

the PA200 or PA28, the 20S proteasome mediates ubiquitin-independent protein degradation. This type of proteolysis is required in several pathways including

spermatogenesis (20S-PA200 complex) or generation of a subset of MHC class I-presented

antigenic peptides (20S-PA28 complex).[UniProtKB/Swiss-Prot Function]