

## **Product datasheet for MR228308**

## Xaf1 (NM\_001291153) Mouse Tagged ORF Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Xaf1 (NM\_001291153) Mouse Tagged ORF Clone

Tag: Myc-DDK

Symbol: Xaf1

Synonyms: Fbox39

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

ORF Nucleotide >MR228308 representing NM\_001291153

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

AGTGCCAGAGGTTAGCTCACCAAAAGAAACTCCAGTGGGGCTGG

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >MR228308 representing NM\_001291153

Red=Cloning site Green=Tags(s)

MEADFQVCRNCKRNVASLHFMLHEAHCLRFIVLCPECEEPIPESKMKEHMEVVHQQKQCSAPNTVTRIRD ESIIVIPSTLAFMDSGNRRSTVSKDVRPKTKNRNSSTKRETKKONGTVALPLKSGLQORADLPTGDETAY

DTLQNCCQCRILLPLPILNEHQEKCQRLAHQKKLQWGW

**TRTRPL**EQKLISEEDLAANDILDYKDDDDK**V** 

**Restriction Sites:** Sgfl-Mlul



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM 001291153

ORF Size: 534 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

**RefSeq:** <u>NM 001291153.1</u>, <u>NP 001278082.1</u>

RefSeq Size: 2313 bp
RefSeq ORF: 537 bp
Locus ID: 327959



UniProt ID: Q5NBU8
Cytogenetics: 11 B4
MW: 20.9 kDa

**Gene Summary:** Seems to function as a negative regulator of members of the IAP (inhibitor of apoptosis

protein) family. Inhibits anti-caspase activity of BIRC4. Induces cleavage and inactivation of BIRC4 independent of caspase activation. Mediates TNF-alpha-induced apoptosis and is involved in apoptosis in trophoblast cells. May inhibit BIRC4 indirectly by activating the mitochondrial apoptosis pathway. After translocation to mitochondria, promotes


translocation of BAX to mitochondria and cytochrome c release from mitochondria. Seems to

promote the redistribution of BIRC4 from the cytoplasm to the nucleus, probably

independent of BIRC4 inactivation which seems to occur in the cytoplasm. The BIRC4-XAF1 complex mediates down-regulation of BIRC5/survivin; the process requires the E3 ligase activity of BIRC4. Seems to be involved in cellular sensitivity to the proapoptotic actions of TRAIL. May be a tumor suppressor by mediating apoptosis resistance of cancer cells (By

similarity).[UniProtKB/Swiss-Prot Function]

## **Product images:**



Circular map for MR228308